



Borel Measurability in Linear Algebra

Edward A. Azoff

Proceedings of the American Mathematical Society, Volume 42, Issue 2 (Feb., 1974),
346-350.

Your use of the JSTOR database indicates your acceptance of JSTOR's Terms and Conditions of Use. A copy of JSTOR's Terms and Conditions of Use is available at <http://www.jstor.org/about/terms.html>, by contacting JSTOR at jstor-info@umich.edu, or by calling JSTOR at (888)388-3574, (734)998-9101 or (FAX) (734)998-9113. No part of a JSTOR transmission may be copied, downloaded, stored, further transmitted, transferred, distributed, altered, or otherwise used, in any form or by any means, except: (1) one stored electronic and one paper copy of any article solely for your personal, non-commercial use, or (2) with prior written permission of JSTOR and the publisher of the article or other text.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

Proceedings of the American Mathematical Society is published by American Mathematical Society. Please contact the publisher for further permissions regarding the use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/ams.html>.

Proceedings of the American Mathematical Society
©1974 American Mathematical Society

JSTOR and the JSTOR logo are trademarks of JSTOR, and are Registered in the U.S. Patent and Trademark Office. For more information on JSTOR contact jstor-info@umich.edu.

©2000 JSTOR

BOREL MEASURABILITY IN LINEAR ALGEBRA

EDWARD A. AZOFF

ABSTRACT. It is shown that the usual processes of linear algebra (e.g., finding Jordan forms, eigenvalues, and eigenvectors) can be carried out in a Borel measurable fashion. These results follow easily from a variant of von Neumann's principle of measurable choice and can be applied to the study of Type I_n von Neumann algebras.

Introduction. Denote by M_n the algebra of $n \times n$ (complex) matrices, considered as operating on the Hilbert space C^n . We equip M_n with its usual (operator) norm, along with the subordinate topological and Borel structures.

The main purpose of this paper is to show that many of the processes of linear algebra (e.g., finding Jordan canonical forms) are Borel measurable. The proofs, which turn out to be remarkably simple, depend only on a 'principle of measurable choice' presented as Theorem 1.

The considerations of the preceding paragraph turn out to be important in the study of Type I_n von Neumann algebras. (This is because such algebras can always be represented as algebras of matrices whose entries are continuous functions.) Indeed, upon examining the papers listed in the bibliography, one finds many arguments which are essentially measure-theoretic in nature. The latter half of the present paper shows how such arguments can be replaced by applications of Theorem 1.

Measurable choice in linear algebra. It is easy to see that many of the processes of linear algebra are not continuous. For example, let $A_n = \begin{bmatrix} 0 & 1/n \\ 0 & 0 \end{bmatrix}$. Then the $\{A_n\}_{n=1}^{\infty}$ converge uniformly to the zero matrix but the Jordan canonical form of each A_n is $\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$. Thus the correspondence, $A \rightarrow$ Jordan form of A , is not continuous. Similar examples show that $\#(A)$, the number of distinct eigenvalues of A , does not depend continuously on A , and that there is no continuous way of associating an eigenvector with each matrix.

Received by the editors May 16, 1973.

AMS (MOS) subject classifications (1970). Primary 15A54, 28A20; Secondary 15A21, 47C15.

Key words and phrases. Borel function, linear algebraic process, principle of measurable choice, continuous matrix valued function on a Stonian space.

© American Mathematical Society 1974

The remainder of this section is devoted to showing that the processes of the preceding paragraph can be carried out in a Borel measurable fashion. The main tool in this development is provided by the following principle of measurable choice.

THEOREM 1. *Let X and Y be complete separable metric spaces and E , a closed σ -compact subset of $X \times Y$. Then $\pi_1(E)$ is a Borel set in X and there exists a Borel function $\varphi: \pi_1(E) \rightarrow Y$ whose graph is contained in E .*

PROOF. Let F be a closed subset of Y . Then $E \cap X \times F$ is closed and hence can be expressed as a countable union of compact sets. The natural projection π_1 being continuous, we see that $\pi_1(E \cap X \times F)$ is the countable union of compact sets and hence is Borel. In particular, this applies to $F = Y$, whence $\pi_1(E)$ is Borel.

Fix a dense subset $\{y_n\}_{n=1}^\infty$ in Y . For each $x \in \pi_1(E)$, we define $\varphi_k(x)$ (inductively on k) to be the first y_n satisfying

- (1) E is not disjoint from $\{x\} \times [\text{closed } 1/2^k\text{-ball about } y_n]$, and
- (2) either $k=1$ or the distance between y_n and $\varphi_{k-1}(x)$ is $\leq 1/2^{k-2}$.

An easy induction argument shows the functions $\{\varphi_k\}_{k=1}^\infty$ are all Borel. Thus, their (uniform) limit φ satisfies the conclusion of the theorem. \square

REMARK. Theorem 1 is essentially a 'folk result'; in particular it follows from Theorem 4 (p. 135) and Exercise 18a (p. 149) of [1].

Theorem 1 differs from the usual von Neumann clipping theorem (see e.g., p. 35 of [2]) in that no set of measure zero must be deleted from $\pi_1(E)$.

COROLLARY 2. *There is a Borel function $\psi: M_n \rightarrow M_n$ associating with each matrix A , a unitary matrix $\psi(A)$ such that $\psi(A)^* A \psi(A)$ is upper triangular.*

PROOF. Let $E = \{(A, U, B) \in M_n \times M_n \times M_n \mid U \text{ is unitary, } B \text{ is upper triangular and } U^* A U = B\}$. Clearly E is closed and since any matrix is unitarily equivalent to a matrix in upper triangular form, $\pi_1(E) = M_n$. Apply Theorem 1 with $X = M_n$ and $Y = M_n \times M_n$ to get a Borel function $\varphi: M_n \rightarrow M_n \times M_n$. The proof is completed by setting $\psi = \pi_1 \circ \varphi$. \square

COROLLARY 3. *There is a Borel function $\psi: M_n \rightarrow M_n$ such that for each matrix A , $\psi(A)$ implements the similarity between A and its Jordan canonical form.*

PROOF. Fix an integer k , and set

$$E_k = \{(A, J, P) \in M_n \times M_n \times M_n \mid J \text{ is in Jordan form, } \|P\| \leq k, \|P^{-1}\| \leq k \text{ and } PAP^{-1} = J\}.$$

We now apply Theorem 1 exactly as in the previous corollary to get a Borel function $\psi_k: \pi_1(E_k) \rightarrow \pi_3(E_k)$.

Now $\bigcup_{k=1}^{\infty} \pi_1(E_k) = M_n$ since every matrix has a Jordan form. Thus we take

$$\begin{aligned} \psi(A) &= \psi_1(A) \quad \text{for } A \in \pi_1(E_1), \\ &= \psi_k(A) \quad \text{for } A \in \pi_1(E_k) \setminus \pi_1(E_{k-1}), \quad k > 1, \end{aligned}$$

and the corollary is proved. \square

COROLLARY 4. (1) $\mathcal{J}(A)$, 'the' Jordan canonical form of A , depends measurably and boundedly on A .

(2) Let G_k (respectively H_k) be the set of matrices in M_n which have k eigenvalues repeated (respectively not repeated) according to multiplicity. Then the $\{G_k\}$ and $\{H_k\}$ are all Borel sets.

(3) There exist k Borel functions $\lambda_1(A) \cdots \lambda_k(A): H_k \rightarrow \mathbb{C}$ which together trace out the spectrum of A .

(4) There exist k Borel functions $v_1(A) \cdots v_k(A): G_k \rightarrow \mathbb{C}^n$ such that each $v_i(A)$ is an eigenvector for A and every eigenvector of A is a linear combination of $v_1(A) \cdots v_k(A)$.

PROOF. Let ψ be as in Corollary 3.

(1) $\mathcal{J}(A) = \psi(A)A[\psi(A)]^{-1}$ is a Borel function of A since ψ is Borel. Also $\|\mathcal{J}(A)\| \leq 1 + \sup\{\text{eigenvalues of } A\} \leq 1 + \|A\|$.

(2), (3), (4). The eigenvalues and eigenvectors of A can be read off from $\mathcal{J}(A)$ and $\psi(A)$. \square

Applications to $M_n(X)$. Let (X, μ) be a perfect Stonian space and denote by $M_n(X)$, the C^* -algebra of continuous functions from X to M_n . (The algebraic operations on $M_n(X)$ are defined pointwise and the norm is the sup norm: $\|A\| \equiv \sup_{t \in X} \|A(t)\|$.) The algebras $M_n(X)$ have been widely studied (see references); their importance lies in the fact [7] that (to within unitary equivalence) they represent all homogeneous Type I_n von Neumann algebras.

To say that (X, μ) is perfect Stonian means that $C(X) = L_\infty(X)$, i.e., every equivalence class of bounded measurable functions on X has a unique continuous representative. In particular, a subset S of X has full measure ($\mu(X \setminus S) = 0$) if and only if S is dense in X . These facts are systematically exploited in the study of $M_n(X)$. For example, let $A, B \in M_n(X)$ and consider the problem of finding a $Y \in M_n(X)$ such that $AY = B$. Thus we want 'pointwise solutions' $\{Y(t)\}_{t \in X}$ depending continuously on t . Suppose we could find pointwise solutions $\{\tilde{Y}(t)\}_{t \in X}$ depending boundedly and measurably on t . Then \tilde{Y} has a unique continuous representative Y and $A(t)Y(t) = B(t)$ for almost all t . Since two continuous functions agreeing on a dense subset must coincide, we see that $AY = B$.

Thus we do not have to put solutions together continuously—it suffices to assemble them boundedly and measurably. It is the brunt of Theorem 1 that even the measurability requirement may be dropped.

COROLLARY 5. *Let $A, B \in M_n(X)$ and $K \geq 0$. The following are equivalent:*

- (1) *The equation $AY=B$ has a solution (in $M_n(X)$) of norm $\leq K$.*
- (2) *For almost all $t \in X$, the equation $A(t)Y=B(t)$ has a solution (in M_n) of norm $\leq K$.*

PROOF. Clearly (1) \Rightarrow (2). Thus assume (2). Let

$$E = \{(A, B, Y) \in M_n \times M_n \times M_n \mid AY = B, \|Y\| \leq K\}.$$

Applying Theorem 1 with $X=M_n \times M_n$ and $Y=M_n$, we get a Borel function $\varphi: \pi_{M_n \times M_n}(E) \rightarrow M_n$. Set

$$\begin{aligned} \tilde{Y}(t) &= \varphi(A(t), B(t)), & \text{when this is defined,} \\ &= 0, & \text{otherwise.} \end{aligned}$$

Let Y be the unique continuous representative of \tilde{Y} . Then $A(t)Y(t)=B(t)$ almost everywhere and hence everywhere. \square

The following two results occur as Theorem 2 of [4] and Theorem 1 of [3] respectively. The proofs, being almost identical to that of Corollary 5, are omitted.

COROLLARY 6 (PEARCY). *Let $A \in M_n(X)$. Then there exists a unitary operator $U \in M_n(X)$ such that U^*AU is in upper triangular form.*

COROLLARY 7 (DECKARD-PEARCY). *Let $A, B \in M_n(X)$ and $K \geq 0$. The following are equivalent:*

- (1) *There is an invertible operator $P \in M_n(X)$ satisfying $PAP^{-1}=B$, $\|P\| \leq K$, $\|P^{-1}\| \leq K$.*
- (2) *For almost all t , there is an invertible operator $P(t) \in M_n$ satisfying $P(t)A(t)[P(t)]^{-1}=B(t)$, $\|P(t)\| \leq K$, $\|[P(t)]^{-1}\| \leq K$.*

As a final application of Theorem 1, we give a simple proof of Theorem 4.6 of [7]. This result plays a central role in [5] and [7]. Recall that $R \in M_n(X)$ is said to be *quasi-invertible* if the only $D \in M_n(X)$ satisfying $RD=0$ or $DR=0$ is $D=0$. Two matrices $A, B \in M_n(X)$ are said to be *quasi-similar* if there exist quasi-invertible matrices R and S in $M_n(X)$ satisfying (1) $SA=BS$ and (2) $AR=RB$.

COROLLARY 8 (HOOVER). *Let $A \in M_n(X)$. There exists a $J \in M_n(X)$ such that $J(t)$ is in Jordan canonical form for each t and A is quasi-similar to J .*

PROOF. Let ψ and \mathcal{J} be as in Corollaries 3 and 4 respectively. Set

$$\tilde{S}(t) = \frac{\psi(A(t))}{\|\psi(A(t))\|}, \quad \tilde{R}(t) = \frac{[\psi(A(t))]^{-1}}{\|[\psi(A(t))]^{-1}\|}, \quad \tilde{J}(t) = \mathcal{J}(A(t)),$$

and denote their continuous representatives by S , R , J respectively. We have for almost all t :

- (1) $J(t)$ is in Jordan form,
- (2) $S(t)A(t) = J(t)S(t)$, and
- (3) $A(t)R(t) = R(t)J(t)$.

The matrices involved being continuous, these statements must hold for all t . Moreover, $S(t)$ and $R(t)$ being invertible for almost all t , we see that S and R are quasi-invertible. \square

ADDED IN PROOF. If X is merely assumed to be a Stonian space (i.e., not equipped with a measure) then every Borel function on X differs from a continuous function on a set of first category. Thus Corollaries 5–8 have obvious analogues in this setting.

REFERENCES

General

1. N. Bourbaki, *Eléments de mathématique. I: Les structures fondamentales de l'analyse*. Fasc. VIII. Livre III: *Topologie générale*. Chap. 9: *Utilisation des nombres réels en topologie générale*, 2nd ed., Actualités Sci. Indust., no. 1045, Hermann, Paris, 1958; English transl., Hermann, Paris; Addison-Wesley, Reading, Mass., 1966. MR 30 #3439; 34 #5044b.
2. J. T. Schwartz, *W*-algebras*, Gordon and Breach, New York, 1967. MR 38 #547.

$M_n(X)$

3. D. Deckard and C. Pearcy, *On continuous matrix-valued functions on a Stonian space*, Pacific J. Math. 14 (1964), 857–869. MR 30 #2356.
4. ———, *On matrices over the ring of continuous complex-valued functions on a Stonian space*, Proc. Amer. Math. Soc. 14 (1963), 322–328. MR 26 #5438.
5. N. Dunford, *A spectral theory for certain operators on a direct sum of Hilbert spaces*, Math. Ann. 162 (1965/66), 294–330. MR 32 #8182.
6. H. Gonsior, *Spectral theory for a class of non-normal operators*, Canad. J. Math. 8 (1956), 449–461. MR 18, 915.
7. T. Hoover, *Hyperinvariant subspaces for n -normal operators*, Acta. Sci. Math. (Szeged) 32 (1971), 109–119.
8. H. Radjavi and P. Rosenthal, *Hyperinvariant subspaces for spectral and n -normal operators*, Acta. Sci. Math. (Szeged) 32 (1971), 121–125.

See also references 1, 7, 9, and 10 in [3].

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF GEORGIA, ATHENS, GEORGIA 30601