

RANKS OF TWISTS OF ELLIPTIC CURVES AND HILBERT'S TENTH PROBLEM

B. MAZUR AND K. RUBIN

ABSTRACT. In this paper we investigate the 2-Selmer rank in families of quadratic twists of elliptic curves over arbitrary number fields. We give sufficient conditions on an elliptic curve so that it has twists of arbitrary 2-Selmer rank, and we give lower bounds for the number of twists (with bounded conductor) that have a given 2-Selmer rank. As a consequence, under appropriate hypotheses we can find many twists with trivial Mordell-Weil group, and (assuming the Shafarevich-Tate conjecture) many others with infinite cyclic Mordell-Weil group. Using work of Poonen and Shlapentokh, it follows from our results that if the Shafarevich-Tate conjecture holds, then Hilbert's Tenth Problem has a negative answer over the ring of integers of every number field.

1. INTRODUCTION AND MAIN RESULTS

In this paper we investigate the 2-Selmer rank in families of quadratic twists of elliptic curves over arbitrary number fields. We give sufficient conditions on an elliptic curve so that it has twists of 2-Selmer rank r for every $r \geq 0$, and discuss other conditions under which the 2-Selmer ranks of all quadratic twists have the same parity. We also give lower bounds for the number of twists (with bounded conductor) that have a given 2-Selmer rank.

Since the 2-Selmer rank is an upper bound for the Mordell-Weil rank, our results have consequences for the Mordell-Weil rank. Under appropriate hypotheses we can find many twists with trivial Mordell-Weil group, and (assuming the Shafarevich-Tate conjecture below) many others with infinite cyclic Mordell-Weil group. Combining our results with work of Poonen and Shlapentokh, we show that if the Shafarevich-Tate conjecture holds, then Hilbert's Tenth Problem has a negative answer over the ring of integers of every number field.

If K is a number field and E is an elliptic curve over K , let $\text{Sel}_2(E/K)$ be the 2-Selmer group of E/K (see §2 for the definition) and

$$d_2(E/K) := \dim_{\mathbf{F}_2} \text{Sel}_2(E/K).$$

Then $\text{rank}(E(K)) \leq d_2(E/K)$, so

$$d_2(E/K) = 0 \implies \text{rank}(E(K)) = 0.$$

If F/K is a quadratic extension, let E^F denote the quadratic twist of E by F/K . We will allow the "trivial quadratic extension" $F = K$, in which case $E^F = E$. For $X \in \mathbf{R}^+$ define

$$N_r(E, X) := |\{\text{quadratic } F/K : d_2(E^F/K) = r \text{ and } \mathbf{N}_{K/\mathbf{Q}}\mathfrak{f}(F/K) < X\}|$$

where $\mathfrak{f}(F/K)$ denotes the finite part of the conductor of F/K .

This material is based upon work supported by the National Science Foundation under grants DMS-0700580 and DMS-0757807.

1.1. Controlling the Selmer rank. Not all elliptic curves have twists of every 2-Selmer rank. For example, some elliptic curves have “constant 2-Selmer parity”, meaning that $d_2(E^F/K) \equiv d_2(E/K) \pmod{2}$ for all quadratic extensions F/K . A theorem of T. Dokchitser and V. Dokchitser [DD, Theorem 1] (see Theorem 9.3 below), combined with standard conjectures, predicts that E/K has constant 2-Selmer parity if and only if K is totally imaginary and E acquires everywhere good reduction over an abelian extension of K . See §9 for a discussion of the phenomenon of constant 2-Selmer parity, and some examples.

We expect that constant parity and the existence of rational 2-torsion are the only obstructions to having twists of every 2-Selmer rank. We also expect that $N_r(E, X)$ should grow like a positive constant times X , whenever it is nonzero. Namely, we expect the following.

Conjecture 1.1. *Suppose K is a number field and E is an elliptic curve over K .*

- (i) *If $r \geq \dim_2 E(K)[2]$ and $r \equiv d_2(E/K) \pmod{2}$, then $N_r(E, X) \gg X$.*
- (ii) *If K has a real embedding, or if E/K does not acquire everywhere good reduction over an abelian extension of K , then $N_r(E, X) \gg X$ for every $r \geq \dim_{\mathbf{F}_2} E(K)[2]$.*

When $K = \mathbf{Q}$ and E is $y^2 = x^3 - x$, Heath-Brown [HB] has shown that $\lim_{X \rightarrow \infty} N_r(E, X)/X = \alpha_r$ for every $r \geq 2$, with an explicit positive constant α_r . Related results have been obtained by Swinnerton-Dyer [SD] when $K = \mathbf{Q}$ and E is an elliptic curve with all 2-torsion points rational.

In the direction of Conjecture 1.1, we have the following results.

Theorem 1.2. *Suppose K is a number field, E is an elliptic curve over K , $r \geq 0$, and E has a quadratic twist E'/K with $d_2(E'/K) = r$. Then:*

- (i) *If $\text{Gal}(K(E[2])/K) \cong S_3$, then $N_r(E, X) \gg X/(\log X)^{2/3}$.*
- (ii) *If $\text{Gal}(K(E[2])/K) \cong \mathbf{Z}/3\mathbf{Z}$, then $N_r(E, X) \gg X/(\log X)^{1/3}$.*

Note that $\text{Gal}(K(E[2])/K)$ is isomorphic to S_3 or $\mathbf{Z}/3\mathbf{Z}$ if and only if $E(K)[2] = 0$.

When $K = \mathbf{Q}$, a version of Theorem 1.2 was proved by Chang in [Ch1, Theorem 4.10]. Also in the case $K = \mathbf{Q}$, Chang has proved (slightly weaker) versions of Theorem 1.5 and Corollary 1.10 below, namely [Ch2, Theorem 1.1] and [Ch2, Corollary 1.2], respectively.

In the statements below, we will use the phrase “ E has many twists” with some property to indicate that the number of such twists, ordered by $\mathbf{N}_{K/\mathbf{Q}}\mathfrak{f}(F/K)$, is $\gg X/(\log X)^\alpha$ for some $\alpha \in \mathbf{R}$.

Theorem 1.3. *Suppose K is a number field, and E is an elliptic curve over K such that $E(K)[2] = 0$. Suppose further that either K has a real embedding, or that E has multiplicative reduction at some prime of K .*

If $0 \leq r \leq \max\{d_2(E/K), 1\}$, then E has many quadratic twists E'/K with $d_2(E'/K) = r$.

Theorem 1.4. *Suppose K is a number field, and E is an elliptic curve over K such that $\text{Gal}(K(E[2])/K) \cong S_3$. Let Δ_E be the discriminant of some model of E , and suppose further that K has a place v_0 satisfying one of the following conditions:*

- v_0 is real and $(\Delta_E)_{v_0} < 0$, or
- $v_0 \nmid 2\infty$, E has multiplicative reduction at v_0 , and $\text{ord}_{v_0}(\Delta_E)$ is odd.

Then for every $r \geq 0$, E has many quadratic twists E'/K with $d_2(E'/K) = r$.

Theorem 1.5. *Suppose K is a number field, and E is an elliptic curve over K such that $E(K)[2] = 0$. If $0 \leq r \leq d_2(E/K)$ and $r \equiv d_2(E/K) \pmod{2}$, then E has many quadratic twists E'/K such that $d_2(E'/K) = r$.*

Corollary 1.6. *Suppose K is a number field, and E is an elliptic curve over K with constant 2-Selmer parity such that $\text{Gal}(K(E[2])/K) \cong S_3$. Let $j(E)$ be the j -invariant of E , and suppose further that $j(E) \neq 0$ and K has an archimedean place v such that $(j(E))_v \in \mathbf{R}$ and $(j(E))_v < 1728$. Then for every $r \equiv d_2(E/K) \pmod{2}$, E has many quadratic twists E'/K such that $d_2(E'/K) = r$.*

For every number field K , there are elliptic curves E over K satisfying the hypotheses of Theorem 1.4. In fact, E can be taken to be the base change of an elliptic curve over \mathbf{Q} (see Lemma 5.4).

Corollary 1.7. *Suppose K is a number field. There are elliptic curves E over K such that for every $r \geq 0$, E has many twists E'/K with $d_2(E'/K) = r$.*

1.2. Controlling the Mordell-Weil rank. Using the relation between $d_2(E/K)$ and $\text{rank}(E(K))$ leads to the following corollaries.

Corollary 1.8. *Suppose K is a number field, and E is an elliptic curve over K such that $E(K)[2] = 0$. Suppose further that either K has a real embedding, or that E has multiplicative reduction at some prime of K . Then E has many twists E'/K with $E'(K) = 0$.*

When $K = \mathbf{Q}$, Corollary 1.8 was proved by Ono and Skinner ([OS, §1], [On, Corollary 3]), using methods very different from ours (modularity and special values of L -functions).

Corollary 1.9. *Suppose K is a number field. There are elliptic curves E over K such that E has many twists E'/K with $E'(K) = 0$.*

If E is an elliptic curve over a number field K , let $\text{III}(E/K)$ denote the Shafarevich-Tate group of E over K (see §2). A conjecture that is part of the folklore (usually called the Shafarevich-Tate Conjecture [Ca1, p. 239, footnote (5)]) predicts that $\text{III}(E/K)$ is finite. If the 2-primary subgroup $\text{III}(E/K)[2^\infty]$ is finite, then the Cassels pairing shows that $\dim_{\mathbf{F}_2} \text{III}(E/K)[2]$ is even. We record this 2-parity conjecture as follows.

Conjecture $\text{III}T_2(K)$. *For every elliptic curve E/K , $\dim_{\mathbf{F}_2} \text{III}(E/K)[2]$ is even.*

Corollary 1.10. *Suppose K is a number field, and E is an elliptic curve over K such that $E(K)[2] = 0$. Suppose further that either K has a real embedding, or that E has multiplicative reduction at some prime of K . If Conjecture $\text{III}T_2(K)$ holds, then E has many quadratic twists with infinite cyclic Mordell-Weil group.*

1.3. Controlling the rank over two fields simultaneously. Suppose L/K is a cyclic extension of prime degree of number fields. With care, we can simultaneously control the 2-Selmer rank of twists of E over K and over L , leading to the following result.

Theorem 1.11. *Suppose L/K is a cyclic extension of prime degree of number fields. Then there is an elliptic curve E over K with $\text{rank}(E(L)) = \text{rank}(E(K))$.*

If Conjecture $\text{III}T_2(K)$ is true, then there is an elliptic curve E over K with $\text{rank}(E(L)) = \text{rank}(E(K)) = 1$.

By using the final assertion of Lemma 5.4 in the proof of Theorem 1.11, we can take the elliptic curve E in Theorem 1.11 to be a twist over K of an elliptic curve defined over \mathbf{Q} . Similarly, in Corollaries 1.7 and 1.9 we can conclude that there are elliptic curves E/\mathbf{Q} that have many quadratic twists E'/K having $d_2(E'/K) = r$ or $E'(K) = 0$, respectively.

Using ideas of Poonen and Shlapentokh ([Po, Theorem 1 and Corollary 2], [De], [Sh]), Theorem 1.11 has the following application to Hilbert's Tenth Problem.

Theorem 1.12. *Suppose Conjecture III $T_2(K)$ holds for every number field K . Then for every number field K , Hilbert's Tenth Problem is undecidable (i.e., has a negative answer) over the ring of integers of K .*

1.4. Some remarks about the proofs. Our methods are different from the classical 2-descent, and are more in the spirit of the work of Kolyvagin, especially as described in [MR1]. If F is a quadratic extension of K , the 2-Selmer group $\text{Sel}_2(E^F/K)$ is defined as a subgroup of $H^1(K, E^F[2])$ cut out by local conditions (see Definition 2.1). The G_K -modules $E[2]$ and $E^F[2]$ are canonically isomorphic, so we can view $\text{Sel}_2(E^F/K) \subset H^1(K, E[2])$ for every F . In other words, all the different 2-Selmer groups are subgroups of $H^1(K, E[2])$, cut out by different local conditions. Our method is to try to construct F so that the local conditions defining $\text{Sel}_2(E/K)$ and $\text{Sel}_2(E^F/K)$ agree everywhere except at most one place, and to use that one place to vary the 2-Selmer rank in a controlled manner.

For example, to prove Theorem 1.2 we find many different quadratic extensions F for which *all* of the local conditions defining $\text{Sel}_2(E/K)$ and $\text{Sel}_2(E^F/K)$ are the same, so in fact $\text{Sel}_2(E^F/K) = \text{Sel}_2(E/K)$.

For another example, suppose the hypotheses of Theorem 1.4 are satisfied. We will take $F = \mathbf{Q}(\sqrt{\pi})$, where π is a generator of a prime ideal \mathfrak{p} chosen using the Chebotarev theorem, so that the local conditions defining $\text{Sel}_2(E/K)$ and $\text{Sel}_2(E^F/K)$ are the same for all places different from \mathfrak{p} . By choosing the prime \mathfrak{p} appropriately, we will also ensure that $\text{Sel}_2(E^F/K) \subset \text{Sel}_2(E/K)$ with codimension one, so $d_2(E^F/K) = d_2(E/K) - 1$.

Similarly, we can choose a different F such that $\text{Sel}_2(E/K) \subset \text{Sel}_2(E^F/K)$ with codimension one, so $d_2(E^F/K) = d_2(E/K) + 1$. Now Theorem 1.4 follows by induction.

Theorems 1.3, 1.5, and 1.11 are proved in the same general manner.

A key tool in several of our arguments is a theorem of Kramer [Kr, Theorem 1] that gives a formula for the parity of $d_2(E/K) + d_2(E^F/K)$ in terms of local data. See Theorem 2.3 below.

1.5. Layout of the paper. In the next section we define the 2-Selmer group and study the local subgroups that occur in the definition. In §3 we give a general result (Proposition 3.3) comparing the 2-Selmer ranks of quadratic twists, and lay the groundwork (Lemma 3.5) for using the Chebotarev theorem to construct useful twists.

Theorem 1.2 is proved in §4. Theorems 1.3, 1.4 and 1.5, and Corollaries 1.6, 1.7, 1.8, 1.9, and 1.10, are all proved in §5. In §6 we prove Theorem 1.11 in the case $[L : K] = 2$, and the rest of Theorem 1.11 is proved in §7. Theorem 1.12 is proved in §8. In §9 we discuss elliptic curves with constant 2-Selmer parity.

1.6. Acknowledgements. The authors would like to thank Bjorn Poonen for asking the questions that led to this work. They also thank Poonen and Alexandra

Shlapentokh for explaining how Theorem 1.11 implies Theorem 1.12, and for allowing us to describe their proof in §8.

2. LOCAL CONDITIONS

Fix for this section a number field K .

Definition 2.1. Suppose E is an elliptic curve over K . For every place v of K , let $H_f^1(K_v, E[2])$ denote the image of the Kummer map

$$E(K_v)/2E(K_v) \longrightarrow H^1(K_v, E[2]).$$

Note that $H_f^1(K_v, E[2])$ depends on E , not just on the Galois module $E[2]$. However, if $v \nmid 2\infty$ and E has good reduction at v , then (see for example [Ca2])

$$H_f^1(K_v, E[2]) = H^1(K_v^{\text{ur}}/K, E[2]) \subset H^1(K_v, E[2])$$

and so

$$(1) \quad H_f^1(K_v, E[2]) \cong E[2]/(\text{Frob}_{\mathfrak{p}} - 1)E[2].$$

with the isomorphism given by evaluating cocycles at the Frobenius automorphism $\text{Frob}_{\mathfrak{p}}$.

The 2-Selmer group $\text{Sel}_2(E/K) \subset H^1(K, E[2])$ is the (finite) \mathbf{F}_2 -vector space defined by the exactness of the sequence

$$0 \longrightarrow \text{Sel}_2(E/K) \longrightarrow H^1(K, E[2]) \longrightarrow \bigoplus_v H^1(K_v, E[2])/H_f^1(K_v, E[2]).$$

The kernel $\text{III}(E/K)[2]$ of multiplication by 2 in the Shafarevich-Tate group of E/K is a (finite) \mathbf{F}_2 -vector space satisfying

$$(2) \quad 0 \longrightarrow E(K)/2E(K) \longrightarrow \text{Sel}_2(E/K) \longrightarrow \text{III}(E/K)[2] \longrightarrow 0.$$

In particular (recall that $d_2(E/K) := \dim_{\mathbf{F}_2} \text{Sel}_2(E/K)$) we have

$$d_2(E/K) = \text{rank}(E(K)) + \dim_{\mathbf{F}_2} E(K)[2] + \dim_{\mathbf{F}_2} \text{III}(E/K)[2],$$

so if Conjecture $\text{III}T_2(K)$ holds, then

$$d_2(E/K) \equiv \text{rank}(E(K)) + \dim_{\mathbf{F}_2} E(K)[2] \pmod{2}.$$

Remark. If E is an elliptic curve over K and E^F is a quadratic twist, then there is a natural identification of Galois modules $E[2] = E^F[2]$. This allows us to view $\text{Sel}_2(E/K), \text{Sel}_2(E^F/K) \subset H^1(K, E[2])$, defined by different sets of local conditions. By choosing F carefully, we can ensure that the local conditions $H_f^1(K_v, E[2]), H_f^1(K_v, E^F[2]) \subset H^1(K_v, E[2])$ coincide for all but at most one v , and then using global duality we will compare $d_2(E/K)$ and $d_2(E^F/K)$.

Fix for the rest of this section an elliptic curve E/K and a quadratic extension F/K . Recall that E^F is the twist of E by F/K . Let Δ_E be the discriminant of some model of E .

Definition 2.2. If v is a place of K , let $NE(K_v) \subset E(K_v)$ denote the image of the norm map $E(F_w) \rightarrow E(K_v)$ (this is independent of the choice of w above v), and define

$$\delta_v(E, F/K) := \dim_{\mathbf{F}_2}(E(K_v)/NE(K_v)).$$

The following theorem of Kramer will play an important role in many of our proofs below.

Theorem 2.3 (Kramer). *We have*

$$d_2(E^F/K) \equiv d_2(E/K) + \sum_v \delta_v(E, F/K) \pmod{2}.$$

Proof. See [Kr, Theorem 1]. A key step in his proof is the following remarkable construction. There are alternating Cassels pairings h_E on $\text{Sel}_2(E/K)$ and h_{E^F} on $\text{Sel}_2(E^F/K)$. Their sum is a new alternating pairing on $\text{Sel}_2(E/K) \cap \text{Sel}_2(E^F/K)$, and Kramer shows [Kr, Theorem 2] that the kernel of $h_E + h_{E^F}$ is $\mathbf{N}_{F/K}\text{Sel}_2(E/F)$. Therefore $\dim_{\mathbf{F}_2}((\text{Sel}_2(E/K) \cap \text{Sel}_2(E^F/K))) \equiv \dim_{\mathbf{F}_2}(\mathbf{N}_{F/K}\text{Sel}_2(E/F)) \pmod{2}$. \square

Lemma 2.4. *Under the identification $H_{\mathbf{f}}^1(K_v, E[2]) = E(K_v)/2E(K_v)$, we have*

$$H_{\mathbf{f}}^1(K_v, E[2]) \cap H_{\mathbf{f}}^1(K_v, E^F[2]) = NE(K_v)/2E(K_v).$$

Proof. This is [Kr, Proposition 7] or [MR2, Proposition 5.2] (the proof given in [MR2] works even if $p = 2$, and even if $v \mid \infty$). \square

Lemma 2.5 (Criteria for equality of local conditions after twist). *If at least one of the following conditions (i)-(v) holds:*

- (i) v splits in F/K , or
- (ii) $v \nmid 2$ and $E(K_v)[2] = 0$, or
- (iii) E has multiplicative reduction at v , F/K is unramified at v , and $\text{ord}_v(\Delta_E)$ is odd, or
- (iv) v is real and $(\Delta_E)_v < 0$, or
- (v) v is a prime where E has good reduction and v is unramified in F/K ,

then $H_{\mathbf{f}}^1(K_v, E[2]) = H_{\mathbf{f}}^1(K_v, E^F[2])$ and $\delta_v(E, F/K) = 0$.

Proof. By Lemma 2.4, we have

$$H_{\mathbf{f}}^1(K_v, E[2]) = H_{\mathbf{f}}^1(K_v, E^F[2]) \iff NE(K_v) = E(K_v) \iff \delta_v(E, F/K) = 0.$$

If v splits in F/K then $NE(K_v) = E(K_v)$.

If $v \nmid 2$ then $E(K_v)$ has a 2-divisible subgroup of finite index. If in addition $E(K_v)[2] = 0$, then $E(K_v)$ is 2-divisible so $E(K_v) = 2E(K_v) \subset NE(K_v)$.

If E has multiplicative reduction at v , F/K is unramified at v , and $\text{ord}_v(\Delta_E)$ is odd, then [Kr, Propositions 1 and 2(a)] shows that $\delta_v(E, F/K) = 0$.

If v is real and $(\Delta_E)_v < 0$, then $E(K_v)$ is connected and $\delta_v(E, F/K) = 0$.

If E has good reduction at v and v is unramified in F/K , then $\delta_v(E, F/K) = 0$ by [Maz, Corollary 4.4]. This completes the proof. \square

Lemma 2.6 (Criterion for transversality of local conditions after twist). *If $v \nmid 2\infty$, E has good reduction at v , and v is ramified in F/K , then*

$$H_{\mathbf{f}}^1(K_v, E[2]) \cap H_{\mathbf{f}}^1(K_v, E^F[2]) = 0, \quad E(K_v)/NE(K_v) \cong H_{\mathbf{f}}^1(K_v, E[2]).$$

Proof. For such v , [Maz, Corollary 4.6] or [MR2, Lemma 5.5] show that $NE(K_v) = 2E(K_v)$. Now the lemma follows from Lemma 2.4. \square

3. COMPARING SELMER GROUPS

We continue to fix a number field K , an elliptic curve E/K , and a quadratic extension F/K .

Definition 3.1. If T is a finite set of places of K , let

$$\text{loc}_T : H^1(K, E[2]) \longrightarrow \bigoplus_{v \in T} H^1(K_v, E[2])$$

denote the sum of the localization maps. Define strict and relaxed 2-Selmer groups $\mathcal{S}_T \subset \mathcal{S}^T \subset H^1(K, E[2])$ by the exactness of

$$\begin{aligned} 0 &\longrightarrow \mathcal{S}^T \longrightarrow H^1(K, E[2]) \longrightarrow \bigoplus_{v \notin T} H^1(K_v, E[2]) / H_f^1(K_v, E[2]), \\ 0 &\longrightarrow \mathcal{S}_T \longrightarrow \mathcal{S}^T \xrightarrow{\text{loc}_T} \bigoplus_{v \in T} H^1(K_v, E[2]). \end{aligned}$$

Then by definition $\mathcal{S}_T \subset \text{Sel}_2(E/K) \subset \mathcal{S}^T$, and we define

$$V_T := \text{loc}_T(\text{Sel}_2(E/K)) \subset \bigoplus_{v \in T} H_f^1(K_v, E[2]).$$

Lemma 3.2. $\dim_{\mathbf{F}_2} \mathcal{S}^T - \dim_{\mathbf{F}_2} \mathcal{S}_T = \sum_{v \in T} \dim_{\mathbf{F}_2} H_f^1(K_v, E[2])$.

Proof. We have exact sequences

$$\begin{aligned} 0 &\longrightarrow \text{Sel}_2(E/K) \longrightarrow \mathcal{S}^T \xrightarrow{\text{loc}_T} \bigoplus_{v \in T} (H^1(K_v, E[2]) / H_f^1(K_v, E[2])) \\ 0 &\longrightarrow \mathcal{S}_T \longrightarrow \text{Sel}_2(E/K) \xrightarrow{\text{loc}_T} \bigoplus_{v \in T} H_f^1(K_v, E[2]). \end{aligned}$$

By Poitou-Tate global duality (see for example [Mi, Theorem I.4.10], [T1, Theorem 3.1], or [Ru, Theorem 1.7.3]), the images of the right-hand maps are orthogonal complements under the (nondegenerate) sum of the local Tate pairings, so their \mathbf{F}_2 -dimensions sum to $\sum_{v \in T} \dim_{\mathbf{F}_2} H_f^1(K_v, E[2])$. The lemma follows directly. \square

Proposition 3.3. *Suppose that all of the following places split in F/K :*

- all primes where E has additive reduction,
- all v of multiplicative reduction such that $\text{ord}_v(\Delta_E)$ is even,
- all primes above 2,
- all real places v with $(\Delta_E)_v > 0$,

and suppose in addition that all v of multiplicative reduction such that $\text{ord}_v(\Delta_E)$ is odd are unramified in F/K .

Let T be the set of (finite) primes \mathfrak{p} of K such that F/K is ramified at \mathfrak{p} and $E(K_{\mathfrak{p}})[2] \neq 0$. Then

$$d_2(E^F/K) = d_2(E/K) - \dim_{\mathbf{F}_2} V_T + d$$

for some d satisfying

$$\begin{aligned} 0 &\leq d \leq \dim_{\mathbf{F}_2} (\bigoplus_{\mathfrak{p} \in T} H_f^1(K_{\mathfrak{p}}, E[2]) / V_T), \\ d &\equiv \dim_{\mathbf{F}_2} (\bigoplus_{\mathfrak{p} \in T} H_f^1(K_{\mathfrak{p}}, E[2]) / V_T) \pmod{2}. \end{aligned}$$

Proof. Let $V_T^F := \text{loc}_T(\text{Sel}_2(E^F/K)) \subset \bigoplus_{\mathfrak{p} \in T} H_f^1(K_{\mathfrak{p}}, E[2])$.

By Lemma 2.5, $H_f^1(K_v, E[2]) = H_f^1(K_v, E^F[2])$ if $v \notin T$. Therefore we have $\mathcal{S}_T \subset \text{Sel}_2(E^F/K) \subset \mathcal{S}^T$, and we have exact sequences

$$\begin{aligned} 0 &\longrightarrow \mathcal{S}_T \longrightarrow \text{Sel}_2(E/K) \xrightarrow{\text{loc}_T} V_T \longrightarrow 0 \\ 0 &\longrightarrow \mathcal{S}_T \longrightarrow \text{Sel}_2(E^F/K) \xrightarrow{\text{loc}_T} V_T^F \longrightarrow 0. \end{aligned}$$

We deduce that

$$(3) \quad d_2(E^F/K) = d_2(E/K) + \dim_{\mathbf{F}_2} V_T^F - \dim_{\mathbf{F}_2} V_T.$$

Let

$$t := \sum_{\mathfrak{p} \in T} \dim_{\mathbf{F}_2} H_f^1(K_{\mathfrak{p}}, E[2]).$$

By Lemma 2.6 we have $\text{Sel}_2(E/K) \cap \text{Sel}_2(E^F/K) = \mathcal{S}_T$, and by the remark above we also have $\text{Sel}_2(E/K) + \text{Sel}_2(E^F/K) \subset \mathcal{S}^T$. Hence

$$(4) \quad \dim_{\mathbf{F}_2} V_T + \dim_{\mathbf{F}_2} V_T^F = \dim_{\mathbf{F}_2}(\text{Sel}_2(E/K)/\mathcal{S}_T) + \dim_{\mathbf{F}_2}(\text{Sel}_2(E^F/K)/\mathcal{S}_T) \\ \leq \dim_{\mathbf{F}_2}(\mathcal{S}^T/\mathcal{S}_T) = t,$$

the final equality by Lemma 3.2.

Recall the local norm index $\delta_v(E, F/K)$ of Definition 2.2. By Lemma 2.5, $\delta_v(E, F/K) = 0$ if $v \notin T$, and by Lemma 2.6,

$$\sum_{\mathfrak{p} \in T} \delta_v(E, F/K) = t,$$

so $d_2(E^F/K) \equiv d_2(E/K) + t \pmod{2}$ by Kramer's congruence (Theorem 2.3). Comparing this with (3) we see that

$$(5) \quad \dim_{\mathbf{F}_2} V_T^F \equiv t - \dim_{\mathbf{F}_2} V_T = \dim_{\mathbf{F}_2}(\oplus_{\mathfrak{p} \in T} H_f^1(K_{\mathfrak{p}}, E[2])/V_T) \pmod{2}.$$

By (4) we have

$$(6) \quad 0 \leq \dim_{\mathbf{F}_2} V_T^F \leq t - \dim_{\mathbf{F}_2} V_T = \dim_{\mathbf{F}_2}(\oplus_{\mathfrak{p} \in T} H_f^1(K_{\mathfrak{p}}, E[2])/V_T).$$

If we let $d = \dim_{\mathbf{F}_2} V_T^F$, then the conclusion of the proposition follows from (3), (5), and (6). \square

Corollary 3.4. *Suppose $E, F/K$, and T are as in Proposition 3.3.*

(i) *If $\dim_{\mathbf{F}_2}(\oplus_{\mathfrak{p} \in T} H_f^1(K_{\mathfrak{p}}, E[2])/V_T) \leq 1$, then*

$$d_2(E^F/K) = d_2(E/K) - 2 \dim_{\mathbf{F}_2} V_T + \sum_{\mathfrak{p} \in T} \dim_{\mathbf{F}_2} H_f^1(K_{\mathfrak{p}}, E[2]).$$

(ii) *If $E(K_{\mathfrak{p}})[2] = 0$ for every $\mathfrak{p} \in T$, then $d_2(E^F/K) = d_2(E/K)$.*

Proof. The first assertion follows directly from Proposition 3.3. For (ii), note that T is empty in this case, so (ii) follows from (i). \square

Let $M := K(E[2])$. If $c \in H^1(K, E[2])$ and $\sigma \in G_K$, let $c(\sigma) \in E[2]/(\sigma - 1)E[2]$ denote the image of c under any cocycle representing c . This is well-defined.

Lemma 3.5. *Suppose $\text{Gal}(M/K) \cong S_3$ and $\sigma \in G_K$. Suppose that C is a finite subgroup of $H^1(K, E[2])$, and $\phi : C \rightarrow E[2]/(\sigma - 1)E[2]$ is a homomorphism.*

Then there is a $\gamma \in G_K$ such that $\gamma|_{MK^{\text{ab}}} = \sigma|_{MK^{\text{ab}}}$ and $c(\gamma) = \phi(c)$ for all $c \in C$.

Proof. Let $\Gamma := \text{Gal}(M/K) \cong \text{Aut}(E[2])$. Then $H^1(\Gamma, E[2]) = 0$, so the restriction map

$$H^1(K, E[2]) \hookrightarrow \text{Hom}(G_M, E[2])^\Gamma$$

is injective.

Fix cocycles $\{c_1, \dots, c_k\}$ representing an \mathbf{F}_2 -basis of C . Then c_1, \dots, c_k restrict to linearly independent homomorphisms $\tilde{c}_1, \dots, \tilde{c}_k \in \text{Hom}(G_M, E[2])^\Gamma$.

Let $N \subset \bar{K}$ be the (abelian) extension of M fixed by $\cap_i \ker(\tilde{c}_i) \subset G_M$. Put $W := G_M / \cap_i \ker(\tilde{c}_i) = \text{Gal}(N/M)$. Then W is an \mathbf{F}_2 -vector space with an action of Γ , $\tilde{c}_1, \dots, \tilde{c}_k$ are linearly independent in $\text{Hom}(W, E[2])^\Gamma$, and

$$(7) \quad \tilde{c}_1 \times \dots \times \tilde{c}_k : W \hookrightarrow E[2]^k$$

is a Γ -equivariant injection. Thus W is isomorphic to a Γ -submodule of the semisimple module $E[2]^k$, so W is also semisimple. But if U is an irreducible constituent of W , then U is also an irreducible constituent of $E[2]^k$, so $U \cong E[2]$. Therefore $W \cong E[2]^j$ for some $j \leq k$. But then $j = \dim_{\mathbf{F}_2} \text{Hom}(W, E[2])^\Gamma \geq k$, so $j = k$ and (7) is an isomorphism.

The group Γ acts trivially on $\text{Gal}((MK^{\text{ab}} \cap N)/M)$, but $\text{Gal}(N/M) = W \cong E[2]^k$ has no nonzero quotients on which Γ acts trivially, so $MK^{\text{ab}} \cap N = M$.

Since (7) is surjective and $MK^{\text{ab}} \cap N = M$, we can choose $\tau \in G_M$ such that $c_i(\tau) = \phi(c_i) - c_i(\sigma)$ for $1 \leq i \leq k$, and $\tau|_{MK^{\text{ab}}} = 1$. Then $c_i(\tau\sigma) = c_i(\tau) + \tau(c_i(\sigma)) = \phi(c_i)$ for every i . Since the c_i represent a basis of C , the proposition is satisfied with $\gamma := \tau\sigma$. \square

Lemma 3.6. *Suppose $E(K)[2] = 0$, and c_1, c_2 are cocycles representing distinct nonzero elements of $H^1(K, E[2])$. Then there is a $\gamma \in G_K$ such that $\gamma|_{MK^{\text{ab}}} = 1$ and $c_1(\gamma), c_2(\gamma)$ are an \mathbf{F}_2 -basis of $E[2]$.*

Proof. Let $\Gamma := \text{Gal}(M/K)$, so either $\Gamma \cong S_3$ or $\Gamma \cong \mathbf{Z}/3\mathbf{Z}$. In either case $E[2]$ is an irreducible Γ -module, and $H^1(\Gamma, E[2]) = 0$, so the restriction map

$$H^1(K, E[2]) \hookrightarrow \text{Hom}(G_M, E[2])^\Gamma$$

is injective. Let \tilde{c}_1, \tilde{c}_2 be the distinct nonzero elements of $\text{Hom}(G_M, E[2])^\Gamma$ obtained by restricting c_1, c_2 to G_M .

For $i = 1, 2$ let N_i be the fixed field of $\ker(\tilde{c}_i)$. Then $\tilde{c}_i : \text{Gal}(N_i/M) \rightarrow E[2]$ is nonzero and Γ -equivariant, so it must be an isomorphism.

Let $N = N_1 \cap N_2$. Since \tilde{c}_i identifies $\text{Gal}(N_i/N)$ with a Γ -stable subgroup of $E[2]$, we either have $N_1 = N_2$ or $N_1 \cap N_2 = M$.

If $N_1 = N_2$, then $\tilde{c}_1, \tilde{c}_2 : \text{Gal}(N/M) \rightarrow E[2]$ are different isomorphisms, so we can find $\tau \in \text{Gal}(N/M)$ such that $\tilde{c}_1(\tau)$ and $\tilde{c}_2(\tau)$ are distinct and nonzero.

If $N_1 \cap N_2 = M$, then again we can find $\tau \in \text{Gal}(N_1 N_2/M)$ such that $\tilde{c}_1(\tau)$ and $\tilde{c}_2(\tau)$ are distinct and nonzero.

Since Γ acts trivially on $\text{Gal}((MK^{\text{ab}} \cap N_1 N_2)/M)$, but $\text{Gal}(N_1 N_2/M) \cong E[2]$ or $E[2]^2$ has no nonzero quotients on which Γ acts trivially, we have $MK^{\text{ab}} \cap N_1 N_2 = M$. Thus we can choose $\gamma \in G_M$ such that $\gamma|_{MK^{\text{ab}}} = 1$ and $\gamma|_{N_1 N_2} = \tau$. This γ has the desired properties. \square

4. PROOF OF THEOREM 1.2

In this section we will prove Theorem 1.2. Suppose K is a number field, N is a finite abelian extension of K , and M is another Galois extension of K .

Fix a nonempty union of conjugacy classes $S \subset \text{Gal}(M/K)$. If \mathfrak{p} is a prime of K unramified in M/K , let $\text{Frob}_{\mathfrak{p}}(M/K)$ denote the Frobenius (conjugacy class) of \mathfrak{p} in $\text{Gal}(M/K)$. Define a set of primes of K

$$\mathcal{P} := \{\mathfrak{p} : \mathfrak{p} \text{ is unramified in } NM/K \text{ and } \text{Frob}_{\mathfrak{p}}(M/K) \subset S\}.$$

and two sets of ideals $\mathcal{N}_1 \subset \mathcal{N}$ of K

$$\begin{aligned}\mathcal{N} &:= \{\mathfrak{a} : \mathfrak{a} \text{ is a squarefree product of primes in } \mathcal{P}\}, \\ \mathcal{N}_1 &:= \{\mathfrak{a} \in \mathcal{N} : [\mathfrak{a}, N/K] = 1\},\end{aligned}$$

where $[\cdot, N/K]$ is the global Artin symbol.

Lemma 4.1. *There is a positive real constant C such that*

$$|\{\mathfrak{a} \in \mathcal{N}_1 : \mathbf{N}_{K/\mathbf{Q}}\mathfrak{a} < X\}| = (C + o(1)) \frac{X}{(\log X)^{1-|S|/[M:K]}}.$$

Proof. The proof is a straightforward adaptation of a result of Serre [Se, Théorème 2.4], who proved this when $K = N = \mathbf{Q}$.

Let $G = \text{Gal}(N/K)$. If $\chi : G \rightarrow \mathbf{C}^\times$ is a character, let

$$f_\chi(s) := \sum_{\mathfrak{a} \in \mathcal{N}} \chi(\mathfrak{a}) \mathbf{N}\mathfrak{a}^{-s} = \prod_{\mathfrak{p} \in \mathcal{P}} (1 + \chi(\mathfrak{p}) \mathbf{N}\mathfrak{p}^{-s})$$

where $\chi(\mathfrak{a}) = \chi([\mathfrak{a}, N/K])$. Then standard methods show that

$$\log f_\chi(s) = \sum_{\mathfrak{p} \in \mathcal{P}} \log(1 + \chi(\mathfrak{p}) \mathbf{N}\mathfrak{p}^{-s}) \sim \sum_{\mathfrak{p} \in \mathcal{P}} \chi(\mathfrak{p}) \mathbf{N}\mathfrak{p}^{-s} \sim \delta_\chi \log\left(\frac{1}{s-1}\right)$$

where

$$\delta_\chi := \begin{cases} 0 & \text{if } \chi \text{ is not the trivial character,} \\ |S|/[M : K] & \text{if } \chi \text{ is trivial,} \end{cases}$$

and we write $g(s) \sim h(s)$ for functions g, h defined on the half-plane $\Re(s) > 1$ to mean that $g(s) - h(s)$ extends to a holomorphic function on $\Re(s) \geq 1$. It follows that

$$\sum_{\mathfrak{a} \in \mathcal{N}_1} \mathbf{N}\mathfrak{a}^{-s} = \frac{1}{[N : K]} \sum_{\chi} f_\chi(s) = (s-1)^{-|S|/[M:K]} g(s)$$

with a function $g(s)$ that is holomorphic and nonzero on $\Re(s) \geq 1$. The lemma now follows from a variant of Ikehara's Tauberian Theorem [Wi, p. 322]. \square

Now fix an elliptic curve E over K with $E[2] = 0$, and let Δ be the discriminant of an integral model of E . Let $N = K(8\Delta\infty)$, the ray class field of K modulo 8Δ and all archimedean places, and let $M := K(E[2])$. Let \mathcal{P} and \mathcal{N}_1 be as defined above, with this N and M and with S the set of elements of order 3 in $\text{Gal}(M/K)$. Since $E(K)[2] = 0$ we have $|S| = 2$.

Proposition 4.2. *Suppose $\mathfrak{a} \in \mathcal{N}_1$. Then there is a quadratic extension F/K of conductor \mathfrak{a} such that $d_2(E^F) = d_2(E)$.*

Proof. Fix $\mathfrak{a} \in \mathcal{N}_1$. Then \mathfrak{a} is principal, with a totally positive generator $\alpha \equiv 1 \pmod{8\Delta}$. Let $F = K(\sqrt{\alpha})$. Then all primes above 2, all primes of bad reduction, and all infinite places split in F/K . If \mathfrak{p} ramifies in F/K then $\mathfrak{p} \mid \mathfrak{a}$, so $\mathfrak{p} \in \mathcal{P}$. Thus the Frobenius of \mathfrak{p} in $\text{Gal}(M/K)$ has order 3, which shows that $E(K_{\mathfrak{p}})[2] = 0$. Now the proposition follows from Corollary 3.4(ii). \square

Proof of Theorem 1.2. Recall that S is the set of elements of order 3 in $\text{Gal}(M/K)$, so

$$\frac{|S|}{[M : K]} = \begin{cases} 1/3 & \text{if } \text{Gal}(M/K) \cong S_3, \\ 2/3 & \text{if } \text{Gal}(M/K) \cong \mathbf{Z}/3\mathbf{Z}. \end{cases}$$

Case 1: $d_2(E/K) = r$. By Proposition 4.2,

$$N_r(E, X) \geq |\{\mathfrak{a} \in \mathcal{N}_1 : \mathbf{N}_{K/\mathbf{Q}}\mathfrak{a} < X\}|.$$

The estimate of Lemma 4.1 for the right-hand side of this inequality proves Theorem 1.2 in this case.

Case 2: $d_2(E/K)$ arbitrary. We have assumed that E has a twist E^L with $d_2(E^L/K) = r$. Every twist $(E^L)^{F'}$ of E^L is also a twist E^F of E , and

$$\mathfrak{f}(F/K) \mid \mathfrak{f}(L/K)\mathfrak{f}(F'/K).$$

so $N_r(E, X) \geq N_r(E^L, X/\mathbf{N}_{K/\mathbf{Q}}\mathfrak{f}(L/K))$. Now Theorem 1.2 for E follows from Theorem 1.2 for E^L , which is proved in Case 1. \square

5. TWISTING TO LOWER AND RAISE THE SELMER RANK

In this section we will use Corollary 3.4 and Lemmas 3.5 and 3.6 to prove Theorems 1.3, 1.4, and 1.5:

- (1) Lemmas 3.5 or 3.6 will provide us with Galois automorphisms that evaluate Selmer cocycles in some useful way,
- (2) the Chebotarev Theorem will provide us with primes whose Frobenius automorphisms are the Galois automorphisms we chose in (1),
- (3) Corollary 3.4 will enable us to calculate $d_2(E^F/K)$, where F is a quadratic extension ramified at one of the primes chosen in (2).

We use Proposition 5.1 below to prove Theorem 1.4, Proposition 5.2 to prove Theorem 1.5, and Proposition 5.3 to prove Theorem 1.3. We also prove Corollaries 1.6, 1.7, 1.8, 1.9, and 1.10.

Proposition 5.1. *Suppose E/K satisfies the hypotheses of Theorem 1.4. Suppose L/K is a quadratic extension (or $L = K$) such that the place v_0 of Theorem 1.4 is unramified in L/K , L'/K is a cyclic extension of odd degree, and Σ is a finite set of places of K .*

- (i) *There is a quadratic twist A of E such that $d_2(A/K) = d_2(E/K) + 1$ and $d_2(A^L/K) = d_2(E^L/K) + 1$.*
- (ii) *If $d_2(E/K) > 0$ and $d_2(E^L/K) > 0$, then there is a quadratic twist A of E such that $d_2(A/K) = d_2(E/K) - 1$ and $d_2(A^L/K) = d_2(E^L/K) - 1$.*
- (iii) *If $\text{Sel}_2(E^L/K) \not\subset \text{Sel}_2(E/K)$ inside $H^1(K, E[2])$, then there is a quadratic twist A of E such that $d_2(A/K) = d_2(E/K) + 1$ and $d_2(A^L/K) = d_2(E^L/K) - 1$.*

In all three cases we can take $A = E^F$, where the quadratic extension F/K satisfies:

- all places in $\Sigma - \{v_0\}$ split in F/K ,
- F/K ramifies at exactly one prime \mathfrak{p} , and that prime satisfies $\mathfrak{p} \notin \Sigma$, \mathfrak{p} is inert in L' , and $E(K_{\mathfrak{p}})[2] \cong \mathbf{Z}/2\mathbf{Z}$.

Proof. Let Δ be the discriminant of (some integral model of) E . Let $M := K(E[2]) = K(E^L[2])$, so M is an S_3 -extension of K containing the quadratic extension $K(\sqrt{\Delta})$. Enlarge Σ if necessary so that it includes all infinite places, all primes above 2, and all primes where either E or E^L has bad reduction. Let $v_0 \nmid 2$ be the distinguished place of Theorem 1.4, either real with $\Delta_{v_0} < 0$, or of multiplicative reduction with $\text{ord}_{v_0}(\Delta)$ odd.

Let \mathfrak{d} be the (formal) product of all places in $\Sigma - \{v_0\}$. Let $K(8\mathfrak{d})$ denote the ray class field of K modulo $8\mathfrak{d}$, and let $K[8\mathfrak{d}]$ denote the maximal 2-power extension of K in $K(8\mathfrak{d})$. Note that $K(\sqrt{\Delta})/K$ is ramified at v_0 but $K[8\mathfrak{d}]/K$ is not, and $[L' : K]$ is odd, so the fields $K[8\mathfrak{d}], L', M$ are linearly disjoint. Therefore we can fix an element $\sigma \in G_K$ such that

- $\sigma|_M \in \text{Gal}(M/K) \cong S_3$ has order 2,
- $\sigma|_{K[8\mathfrak{d}]} = 1$,
- $\sigma|_{L'}$ is a generator of $\text{Gal}(L'/K)$.

It follows in particular that $E[2]/(\sigma - 1)E[2] \cong \mathbf{Z}/2\mathbf{Z}$.

Let $C = \text{Sel}_2(E/K) + \text{Sel}_2(E^L/K) \subset H^1(K, E[2])$, and suppose $\phi : C \rightarrow E[2]/(\sigma - 1)E[2]$ is a homomorphism. By Lemma 3.5 we can find $\gamma \in G_K$ such that

- $\gamma|_{ML'K[8\mathfrak{d}]} = \sigma$,
- $c(\gamma) = \phi(c)$ for every $c \in C$.

Let N be a Galois extension of K containing $ML'K[8\mathfrak{d}]$, large enough so that the restriction of C to N is zero. (For example, one can take the compositum of $L'K(8\mathfrak{d})$ with the fixed field of the intersection of the kernels of the restrictions of $c \in C \hookrightarrow \text{Hom}(G_M, E[2])$.) Let \mathfrak{p} be a prime of K not in Σ , whose Frobenius in $\text{Gal}(N/K)$ is the conjugacy class of γ . Since $\gamma|_{K[8\mathfrak{d}]} = \sigma|_{K[8\mathfrak{d}]} = 1$, and $[K(8\mathfrak{d}) : K[8\mathfrak{d}]]$ is odd, there is an odd positive integer h such that $\gamma^h|_{K(8\mathfrak{d})} = 1$. Thus \mathfrak{p}^h is principal, with a generator $\pi \equiv 1 \pmod{8\mathfrak{d}}$, positive at all real embeddings different from v_0 . Let $F = K(\sqrt{\pi})$. Then all places $v \in \Sigma - \{v_0\}$ split in F , F/K is ramified at \mathfrak{p} and nowhere else, \mathfrak{p} is inert in L'/K because $\gamma|_{L'}$ generates $\text{Gal}(L'/K)$, and $E(K_{\mathfrak{p}})[2] \neq 0$ because $\text{Frob}_{\mathfrak{p}}|_{E[2]} = \sigma|_{E[2]}$ has order 2.

We will apply Corollary 3.4, with $T = \{\mathfrak{p}\}$. Since E has good reduction at \mathfrak{p} , it follows from (1) that

$$(8) \quad H_{\mathfrak{f}}^1(K_{\mathfrak{p}}, E[2]) \cong E[2]/(\text{Frob}_{\mathfrak{p}} - 1)E[2] = E[2]/(\sigma - 1)E[2],$$

and similarly with E replaced by E^L , so

$$\dim_{\mathbf{F}_2} H_{\mathfrak{f}}^1(K_{\mathfrak{p}}, E[2]) = \dim_{\mathbf{F}_2} H_{\mathfrak{f}}^1(K_{\mathfrak{p}}, E^L[2]) = 1.$$

Further, the localization maps

$$\text{loc}_T : \text{Sel}_2(E/K), \text{Sel}_2(E^L/K) \longrightarrow H_{\mathfrak{f}}^1(K_{\mathfrak{p}}, E[2]) \xrightarrow{\sim} E[2]/(\sigma - 1)E[2]$$

are given by evaluation of cocycles at $\text{Frob}_{\mathfrak{p}} = \gamma$. Hence by our choice of γ , (8) identifies

$$\text{loc}_T(\text{Sel}_2(E/K)) = \phi(\text{Sel}_2(E/K)), \quad \text{loc}_T(\text{Sel}_2(E^L/K)) = \phi(\text{Sel}_2(E^L/K)).$$

Thus by Corollary 3.4(i) we conclude that

$$d_2(E^F/K) = \begin{cases} d_2(E/K) + 1 & \text{if } \phi(\text{Sel}_2(E/K)) = 0, \\ d_2(E/K) - 1 & \text{if } \phi(\text{Sel}_2(E/K)) \neq 0. \end{cases}$$

$$d_2((E^F)^L/K) = d_2((E^L)^F/K) = \begin{cases} d_2(E^L/K) + 1 & \text{if } \phi(\text{Sel}_2(E^L/K)) = 0, \\ d_2(E^L/K) - 1 & \text{if } \phi(\text{Sel}_2(E^L/K)) \neq 0. \end{cases}$$

For assertion (i), let $\phi = 0$. For (ii), if $d_2(E/K) > 0$ and $d_2(E^L/K) > 0$, then we can choose a ϕ that is nonzero on both $\text{Sel}_2(E/K)$ and $\text{Sel}_2(E^L/K)$. For (iii), if $\text{Sel}_2(E^L/K) \not\subset \text{Sel}_2(E/K)$, then we can choose a ϕ that is zero on $\text{Sel}_2(E/K)$ and nonzero on $\text{Sel}_2(E^L/K)$. In all three cases, the proposition holds with $A = E^F$. \square

Proof of Theorem 1.4. Note that if E satisfies the hypotheses of Theorem 1.4, then so does every quadratic twist of E .

If $r \geq d_2(E/K)$, then applying Proposition 5.1(i) $r - d_2(E/K)$ times (with $L = L' = K$) shows that E has a twist E' with $d_2(E'/K) = r$.

If $0 \leq r \leq d_2(E/K)$ then applying Proposition 5.1(ii) $d_2(E/K) - r$ times shows that E has a twist E' with $d_2(E'/K) = r$.

Now Theorem 1.2 shows that for every $r \geq 0$, E has many twists E' with $d_2(E'/K) = r$. \square

Proposition 5.2. *Suppose E/K is an elliptic curve such that $E(K)[2] = 0$. If $d_2(E/K) > 1$, then E has a quadratic twist E^F over K such that $d_2(E^F/K) = d_2(E/K) - 2$.*

Proof. The proof is similar to that of Proposition 5.1(ii). Let $M := K(E[2])$, and let Δ be the discriminant of (some integral model of) E . Let $K(8\Delta\infty)$ denote the ray class field of K modulo the product of 8Δ and all infinite places.

Since $d_2(E/K) > 1$, we can choose cocycles c_1, c_2 representing \mathbf{F}_2 -independent elements of $\text{Sel}_2(E/K)$. By Lemma 3.6 we can find $\gamma \in G_K$ such that

- $\gamma|_{MK(8\Delta\infty)} = 1$,
- $c_1(\gamma), c_2(\gamma)$ are an \mathbf{F}_2 -basis of $E[2]$.

Let N be a Galois extension of K containing $MK(8\Delta\infty)$, large enough so that the restriction of $\text{Sel}_2(E/K)$ to N is zero. Let \mathfrak{p} be a prime of K where E has good reduction, not dividing 2, whose Frobenius in $\text{Gal}(N/K)$ is the conjugacy class of γ . Then \mathfrak{p} has a totally positive generator $\pi \equiv 1 \pmod{8\Delta}$. Let $F = K(\sqrt{\pi})$. Then all places v dividing $2\Delta\infty$ split in F/K , and \mathfrak{p} is the only prime that ramifies in F/K .

We will apply Corollary 3.4 with $T = \{\mathfrak{p}\}$. Since E has good reduction at \mathfrak{p} , it follows from (1) that

$$H_{\mathfrak{f}}^1(K_{\mathfrak{p}}, E[2]) = E[2]/(\text{Frob}_{\mathfrak{p}} - 1)E[2] = E[2]/(\gamma - 1)E[2] = E[2].$$

The localization map $\text{loc}_T : \text{Sel}_2(E/K) \rightarrow H_{\mathfrak{f}}^1(K_{\mathfrak{p}}, E[2])$ is given by evaluation of cocycles at $\text{Frob}_{\mathfrak{p}} = \gamma$, so by our choice of γ , the classes $\text{loc}_T(c_1)$ and $\text{loc}_T(c_2)$ generate $H_{\mathfrak{f}}^1(K_{\mathfrak{p}}, E[2])$. In particular loc_T is surjective, so in the notation of Corollary 3.4 we have $\dim_{\mathbf{F}_2} V_T = \dim_{\mathbf{F}_2} H_{\mathfrak{f}}^1(K_{\mathfrak{p}}, E[2]) = 2$. Corollary 3.4(i) now yields $d_2(E^F/K) = d_2(E/K) - 2$, as desired. \square

Proof of Theorem 1.5. Suppose $0 \leq r \leq d_2(E/K)$. Applying Proposition 5.2 $(d_2(E/K) - r)/2$ times shows that E has a twist E' with $d_2(E'/K) = r$, and then Theorem 1.2 shows that E has many such twists. \square

Proof of Corollary 1.6. Let $k = \mathbf{Q}(j(E)) \subset K$. Fix an elliptic curve E_0 over k with $j(E_0) = j(E)$. Since $j(E) \neq 0, 1728$, E_0 is a quadratic twist of E over K . Thus $[k(E_0[2]) : k] \geq [K(E_0[2]) : K] = [K(E[2]) : K]$, so $\text{Gal}(k(E_0[2])/k) \cong S_3$. Also

$$j(E) - 1728 = j(E_0) - 1728 = c_6(E_0)^2/\Delta_{E_0}$$

so $(\Delta_{E_0})_v < 0$ at the real embedding v of k . Therefore E_0/k satisfies the hypotheses of Theorem 1.4, so Theorem 1.4 shows that $d_2(E_0^F/k)$ can be arbitrarily large as F varies through quadratic extensions of k . Since $E(K)[2] = 0$, the map $\text{Sel}_2(E_0^F/k) \rightarrow \text{Sel}_2(E_0^F/K)$ is injective, and so $d_2(E^F/K)$ can be arbitrarily large as F varies through quadratic extensions of K . Now the corollary follows from Theorem 1.5. \square

Proposition 5.3. *Suppose E/K is an elliptic curve such that $E(K)[2] = 0$, and either K has a real embedding, or E has multiplicative reduction at some prime of K . Then E has a quadratic twist E^F/K such that $d_2(E^F/K) \not\equiv d_2(E/K) \pmod{2}$ and $d_2(E^F/K) \geq d_2(E/K) - 1$.*

Proof. Let $M := K(E[2])$, and let Δ be the discriminant of (some integral model of) E . Let \mathfrak{d} be the (formal) product of Δ and all infinite places, let $K(8\mathfrak{d})$ denote the ray class field of K modulo $8\mathfrak{d}$, and let $K[8\mathfrak{d}]$ denote the maximal 2-power extension of K in $K(8\mathfrak{d})$. We have $M \cap K[8\mathfrak{d}] = K(\sqrt{\Delta})$.

Let v_0 be the distinguished place, either real or of multiplicative reduction. Let $\mathbf{x} = (x_v)$ be an idele of K defined by:

- $x_v = 1$ if $v \neq v_0$,
- $x_{v_0} = -1$ if v_0 is real, x_{v_0} is a unit at v_0 such that $K_{v_0}(\sqrt{x_{v_0}})$ is the unramified quadratic extension of K_{v_0} if v_0 is nonarchimedean.

Let $\sigma = [\mathbf{x}, K[8\mathfrak{d}]/K] \in \text{Gal}(K[8\mathfrak{d}]/K)$ be the image of \mathbf{x} under the global Artin map. We consider two cases.

Case 1: $\sigma(\sqrt{\Delta}) = \sqrt{\Delta}$. In this case we can choose $\gamma \in \text{Gal}(MK[8\mathfrak{d}]/K)$ such that $\gamma|_{K[8\mathfrak{d}]} = \sigma$ and $\gamma|_M$ has order 3.

Case 2: $\sigma(\sqrt{\Delta}) = -\sqrt{\Delta}$. In this case $\text{Gal}(M/K) \cong S_3$, and σ is nontrivial on $M \cap K[8\mathfrak{d}] = K(\sqrt{\Delta})$. By Lemma 3.5 we can find $\gamma \in G_K$ such that $\gamma|_{K[8\mathfrak{d}]} = \sigma$, $\gamma|_M$ has order 2, and $c(\gamma) \in (\gamma - 1)E[2]$ for every cocycle c representing an element of $\text{Sel}_2(E/K)$.

In either case, let \mathfrak{p} be a prime of K not dividing 2Δ , whose Frobenius in $\text{Gal}(MK[8\mathfrak{d}]/K)$ is γ . Then some odd power \mathfrak{p}^h is principal, with a generator π such that $\pi \in (K_v^\times)^2$ if $v \mid 2\Delta\infty$ and $v \neq v_0$, $K_{v_0}(\sqrt{\pi}) = \mathbf{C}$ if v_0 is real, and $K_{v_0}(\sqrt{\pi})$ is the unramified quadratic extension of K_{v_0} if v_0 is nonarchimedean.

Let $F = K(\sqrt{\pi})$, and recall the local norm index $\delta_v(E, F/K)$ of Definition 2.2. All places $v \mid 2\Delta\infty$ different from v_0 split in F/K , so by Lemma 2.5, $\delta_v(E, F/K) = 0$ and $H_f^1(K_v, E[2]) = H_f^1(K_v, E^F[2])$ if $v \neq v_0, \mathfrak{p}$. It follows (using Kramer's congruence Theorem 2.3 for (9)) that

$$(9) \quad d_2(E^F/K) \equiv d_2(E/K) + \delta_{v_0}(E, F/K) + \delta_{\mathfrak{p}}(E, F/K) \pmod{2},$$

and

$$(10) \quad \ker[\text{Sel}_2(E/K) \longrightarrow H_f^1(K_{v_0}, E[2]) \oplus H_f^1(K_{\mathfrak{p}}, E[2])] \subset \text{Sel}_2(E^F/K).$$

Consider the Hilbert symbol $(\Delta, \pi)_v$, which is 1 if Δ is a norm from $(F \otimes K_v)^\times$ to K_v^\times , and -1 if not. Then $(\Delta, \pi)_v = 1$ if $v \neq v_0, \mathfrak{p}$, and $\prod_v (\Delta, \pi)_v = 1$, so $(\Delta, \pi)_{v_0} = (\Delta, \pi)_{\mathfrak{p}}$. By [Kr, Proposition 6] if v_0 is real, and by [Kr, Propositions 1, 2] if v_0 is multiplicative, we have

$$\delta_{v_0}(E, F/K) = \begin{cases} 1 & \text{if } (\Delta, \pi)_{v_0} = 1 \\ 0 & \text{if } (\Delta, \pi)_{v_0} = -1. \end{cases}$$

By [Kr, Proposition 3], and using that γ acts nontrivially on $E[2]$ in both Case 1 and Case 2, we have

$$\delta_{\mathfrak{p}}(E, F/K) = \begin{cases} 0 & \text{if } (\Delta, \pi)_{\mathfrak{p}} = 1 \\ 1 & \text{if } (\Delta, \pi)_{\mathfrak{p}} = -1. \end{cases}$$

Thus $\delta_{v_0}(E, F/K) + \delta_{\mathfrak{p}}(E, F/K) = 1$, so (9) shows that $d_2(E^F/K)$ and $d_2(E/K)$ have opposite parity.

In Case 1, $E[2]/(\gamma - 1)E[2] = 0$, so $H_{\mathfrak{f}}^1(K_{\mathfrak{p}}, E[2]) = 0$ by (1). In Case 2, the restriction map $\text{Sel}_2(E/K) \rightarrow H_{\mathfrak{f}}^1(K_{\mathfrak{p}}, E[2]) \cong E[2]/(\gamma - 1)E[2]$ is given by evaluating cocycles at γ , so by our choice of γ this image is zero. In both cases, $\dim_{\mathbf{F}_2} H_{\mathfrak{f}}^1(K_{v_0}, E[2]) \leq 2$, so by (10) we have $d_2(E^F/K) \geq d_2(E/K) - 2$. This completes the proof. \square

Proof of Theorem 1.3. Let E^F be a twist of E as in Proposition 5.3. Theorem 1.3 follows directly from Theorem 1.5 applied to E and to E^F . \square

Lemma 5.4. *Suppose \mathfrak{p} is a prime of K not dividing 2. Then there is an elliptic curve E/K with all of the following properties:*

- (i) E is semistable at all primes,
- (ii) E has multiplicative reduction at \mathfrak{p} and $\text{ord}_{\mathfrak{p}}(\Delta_E) = 1$,
- (iii) $\text{Gal}(K(E[2])/K) \cong S_3$.

If in addition the rational prime p below \mathfrak{p} is unramified in the Galois closure of K/\mathbf{Q} , then E can be taken to be the base change of an elliptic curve over \mathbf{Q} .

Proof. Let E_t be the elliptic curve $y^2 + y = x^3 - x^2 + t$ over $K(t)$. Then

$$j(E_t) = -\frac{2^{12}}{(4t+1)(108t+11)}, \quad \Delta(E_t) = (4t+1)(108t+11), \quad c_4(E_t) = 16.$$

It follows from [S1, Proposition VII.5.1] that for every $t \in \mathcal{O}_K$, E_t has semistable reduction at all primes of K .

Let $\eta \in \mathcal{O}_K$ be such that $\text{ord}_{\mathfrak{p}}(4\eta + 1) = 1$, and let $g(t) := \eta + (4\eta + 1)^2 t$. Then for every $t \in \mathcal{O}_K$ we have $\text{ord}_{\mathfrak{p}}(4g(t) + 1) = 1$. The splitting field of $f_t(x) := x^3 - x^2 + g(t) + 1/4$ over $K(t)$ has Galois group S_3 , since f_t is irreducible and its discriminant $-4(2g(t) + 1)(54g(t) + 19)$ is not a square. Hence by Hilbert's Irreducibility Theorem, there is an integer $t_0 \in \mathcal{O}_K$ such that the splitting field of $f_{t_0}(x)$ over K is an S_3 -extension.

Let E be the elliptic curve $E_{g(t_0)}$. Then $K(E[2])$ is the splitting field of $f_{t_0}(x)$, so $\text{Gal}(K(E[2])/K) \cong S_3$, and

$$\Delta(E) = (4g(t) + 1)(108g(t) + 11) = (4g(t) + 1)(27(4g(t) + 1) - 16)$$

Thus E satisfies (i), (ii), and (iii).

Let K' be the Galois closure of K/\mathbf{Q} , and p the rational prime below \mathfrak{p} , and suppose p is unramified in K'/\mathbf{Q} . We can apply the lemma with p and \mathbf{Q} in place of \mathfrak{p} and K to produce a semistable elliptic curve E/\mathbf{Q} such that $\text{ord}_p(\Delta_E) = 1$ and $\text{Gal}(\mathbf{Q}(E[2])/\mathbf{Q}) \cong S_3$.

Then E/K satisfies (i) and (ii). Further, $\mathbf{Q}(E[2]) \cap K'$ is a Galois extension of \mathbf{Q} that does not contain $\mathbf{Q}(\sqrt{\Delta_E})$ (since the latter is ramified at p). Therefore $\mathbf{Q}(E[2]) \cap K = \mathbf{Q}$, and so $\text{Gal}(K(E[2])/K) \cong \text{Gal}(\mathbf{Q}(E[2])/\mathbf{Q}) \cong S_3$. \square

Proof of Corollary 1.7. By Lemma 5.4, we can find an elliptic curve E over K and a prime $\mathfrak{p} \nmid 2$ such that E has multiplicative reduction at \mathfrak{p} , $\text{ord}_{\mathfrak{p}}(\Delta_E) = 1$, and $\text{Gal}(K(E[2])/K) \cong S_3$. By Theorems 1.4 and 1.2, this E has many quadratic twists E' with $d_2(E'/K) = r$, for every $r \geq 0$. \square

Lemma 5.5. *Suppose E is an elliptic curve over K . Then for all but finitely many quadratic twists E' of E , $E'(K)$ has no odd-order torsion.*

Proof. This is proved in [GM, Proposition 1] when $K = \mathbf{Q}$; we adapt the proof given there. By Merel's Uniform Boundedness Theorem for torsion on elliptic curves [Me], the set

$$\{\text{primes } p : E^F(K)[p] \neq 0 \text{ for some quadratic extension } F/K\}$$

is finite. On the other hand, if p is odd and $\rho_p : G_K \rightarrow \text{Aut}(E[p]) \cong \text{GL}_2(\mathbf{F}_p)$ denotes the mod- p representation attached to E , then there are at most two characters χ of G_K such that $\rho_p \otimes \chi$ contains a copy of the trivial representation. Therefore for fixed odd p , the set

$$\{F/K \text{ quadratic} : E^F(K)[p] \neq 0\}$$

has order at most 2. This completes the proof. \square

Proof of Corollary 1.8. By Theorems 1.3 and 1.2, E has many quadratic twists E' with $d_2(E'/K) = 0$, and hence $\text{rank}(E'(K)) = 0$ by (2). Since $E(K)[2] = 0$, none of these twists have rational 2-torsion, and by Lemma 5.5, only finitely many of these twists have odd-order torsion. This proves the corollary. \square

Proof of Corollary 1.9. By Lemma 5.4 there is an elliptic curve E over K with multiplicative reduction at a prime $\mathfrak{p} \nmid 2$, and with $E[2] = 0$. Now the corollary follows from Corollary 1.8. \square

Proof of Corollary 1.10. By Theorems 1.3 and 1.2, E has many quadratic twists E' with $d_2(E'/K) = 1$. Since $E(K)[2] = 0$, it follows from (2) that either $\text{rank}(E'(K)) = 1$ or $\dim_{\mathbf{F}_2} \text{III}(E'/K)[2] = 1$. But Conjecture III $T_2(K)$ says that $\dim_{\mathbf{F}_2} \text{III}(E'/K)[2]$ is even, so $\text{rank}(E'(K)) = 1$. By Lemma 5.5, all but finitely many of these twists have $E'(K)_{\text{tors}} = 0$, and this proves the corollary. \square

6. PROOF OF THEOREM 1.11 WHEN $[L : K] = 2$

Proposition 6.1. *Suppose L/K is a quadratic extension. Then there is an elliptic curve E/K such that $\text{Gal}(K(E[2])/K) \cong S_3$ and $d_2(E/K) + d_2(E^L/K)$ is odd.*

Proof. Fix a prime $\mathfrak{p} \nmid 6$ that remains prime in L/K . Using Lemma 5.4, fix an elliptic curve E over K with $\text{Gal}(K(E[2])/K) \cong S_3$, with multiplicative reduction at \mathfrak{p} , and with $\text{ord}_{\mathfrak{p}}(\Delta_E) = 1$.

Recall the local norm index $\delta_v(E, L/K)$ of Definition 2.2. We will find a quadratic twist E' of E such that

$$(11) \quad \delta_v(E', L/K) = \begin{cases} \delta_v(E, L/K) & \text{if } v \neq \mathfrak{p}, \\ \delta_v(E, L/K) + 1 & \text{if } v = \mathfrak{p}. \end{cases}$$

Then by Kramer's congruence (Theorem 2.3) we will have

$$d_2(E'/K) + d_2((E')^L/K) \equiv d_2(E/K) + d_2(E^L/K) + 1 \pmod{2}$$

so either E or E' satisfies the conditions of the Proposition.

Let \mathfrak{d} denote the product of all primes different from \mathfrak{p} where E has bad reduction, and let \mathfrak{f} be the conductor of L/K . Let $K(8\mathfrak{d}\mathfrak{f}\infty)$ denote the ray class field of K modulo $8\mathfrak{d}\mathfrak{f}$ and all archimedean primes. Let $M := K(E[2])$, so M is an S_3 -extension of K containing the quadratic field $K(\sqrt{\Delta_E})$. Since $\text{ord}_{\mathfrak{p}}(\Delta_E) = 1$, $K(\sqrt{\Delta_E})/K$ is ramified at \mathfrak{p} , so $K(\sqrt{\Delta_E}) \not\subset K(8\mathfrak{d}\mathfrak{f}\infty)$ and $M \cap K(8\mathfrak{d}\mathfrak{f}\infty) = K$. Therefore we can choose $\sigma \in \text{Gal}(MK(8\mathfrak{d}\mathfrak{f}\infty)/K)$ such that $\sigma|_{K(8\mathfrak{d}\mathfrak{f}\infty)}$ is in the conjugacy class

of the Frobenius of \mathfrak{p} in $\text{Gal}(K(8\mathfrak{d}f\infty)/K)$, and $\sigma|_M$ has order 3. Choose a prime \mathfrak{q} of K whose Frobenius in $\text{Gal}(MK(8\mathfrak{d}f\infty)K)$ is the conjugacy class of σ^{-1} .

It follows that the global Artin map sends $\mathfrak{p}\mathfrak{q}$ to 1 in $\text{Gal}(K(8\mathfrak{d}f\infty)/K)$, so $\mathfrak{p}\mathfrak{q}$ has a totally positive generator $\pi \equiv 1 \pmod{8\mathfrak{d}f}$. Let $F = K(\sqrt{\pi})$, and let E' be the quadratic twist E^F . We need to show (11).

Since $\pi \equiv 1 \pmod{8\mathfrak{d}f}$ and π is totally positive, all primes v dividing $2\mathfrak{d}f\infty$ split in F/K , so $E' \cong E$ over K_v and $\delta_v(E', L/K) = \delta_v(E, L/K)$.

If $v \nmid 2\mathfrak{d}f\mathfrak{p}\mathfrak{q}\infty$, then E and E' have good reduction at v and L/K is unramified at v , so $\delta_v(E', L/K) = \delta_v(E, L/K) = 0$ by Lemma 2.5(v).

Since the Frobenius of \mathfrak{q} in $\text{Gal}(M/K)$ has order 3, we have $E(K_{\mathfrak{q}})[2] = 0$, so $\delta_{\mathfrak{q}}(E', L/K) = \delta_{\mathfrak{q}}(E, L/K) = 0$ by Lemma 2.5(ii).

It remains only to show that $\delta_{\mathfrak{p}}(E', L/K) = \delta_{\mathfrak{p}}(E, L/K) + 1$. Since E has multiplicative reduction at \mathfrak{p} , $\text{ord}_{\mathfrak{p}}(\Delta_E) = 1$, and $L_{\mathfrak{p}}/K_{\mathfrak{p}}$ is unramified, Lemma 2.5(iii) shows that $\delta_{\mathfrak{p}}(E, L/K) = 0$.

On the other hand, since \mathfrak{p} ramifies in F/K , E' has additive reduction at \mathfrak{p} . In fact, Tate's algorithm ([T2] or [S2, §IV.9.4]) shows that E' has reduction type I_1^* at \mathfrak{p} , and (since L/K is unramified at \mathfrak{p}) we have filtrations

$$E'(K_{\mathfrak{p}}) \supset E'_0(K_{\mathfrak{p}}) \supset E'_1(K_{\mathfrak{p}}), \quad E'(L_{\mathfrak{p}}) \supset E'_0(L_{\mathfrak{p}}) \supset E'_1(L_{\mathfrak{p}})$$

where

- $E'(K_{\mathfrak{p}})/E'_0(K_{\mathfrak{p}}) \cong E'(L_{\mathfrak{p}})/E'_0(L_{\mathfrak{p}})$ are cyclic of order 2 or 4,
- $E'_0(K_{\mathfrak{p}})/E'_1(K_{\mathfrak{p}}) \cong \mathcal{O}_K/\mathfrak{p}$ and $E'_0(L_{\mathfrak{p}})/E'_1(L_{\mathfrak{p}}) \cong \mathcal{O}_L/\mathfrak{p}\mathcal{O}_L$,
- $E'_1(K_{\mathfrak{p}})$ and $E'_1(L_{\mathfrak{p}})$ are pro- p groups where p is the residue characteristic of \mathfrak{p} .

It follows that the map $E'(K_{\mathfrak{p}})/2E'(K_{\mathfrak{p}}) \rightarrow E'(L_{\mathfrak{p}})/2E'(L_{\mathfrak{p}})$ is an isomorphism of groups of order 2, so $\mathbf{N}_{L_{\mathfrak{p}}/K_{\mathfrak{p}}} E'(L_{\mathfrak{p}}) = 2E'(K_{\mathfrak{p}})$ and $\delta_{\mathfrak{p}}(E', L/K) = 1$. This completes the proof of (11), and of the proposition. \square

Theorem 6.2. *Suppose L/K is a quadratic extension of number fields. There is an elliptic curve E over K such that $d_2(E/K) = 0$ and $d_2(E^L/K) = 1$. In particular $\text{rank}(E^L(K)) = \text{rank}(E^L(L))$, and if Conjecture III $T_2(K)$ holds then $\text{rank}(E^L(K)) = \text{rank}(E^L(L)) = 1$.*

Proof. Fix an elliptic curve A over K satisfying the conclusion of Proposition 6.1: $\text{Gal}(K(A[2])/K) \cong S_3$ and $d_2(A/K), d_2(A^L/K)$ have opposite parity.

Now apply Proposition 5.1(ii) repeatedly (with $L' = K$), twisting A until we produce a twist B with either $d_2(B/K) = 0$ or $d_2(B^L/K) = 0$. Switching B and B^L if necessary, we may suppose that $d_2(B/K) = 0$.

Note that $d_2(B/K)$ and $d_2(B^L/K)$ still have opposite parity, so $d_2(B^L/K) \geq 1$. If $d_2(B^L/K) = 1$ we stop. If $d_2(B^L/K) > 1$ we apply Proposition 5.1(iii) and then Proposition 5.1(ii), to obtain a twist C with $d_2(C/K) = 0$ and $d_2(C^L/K) = d_2(B^L/K) - 2$. Continuing in this way we eventually obtain a twist E with $d_2(E/K) = 0$ and $d_2(E^L/K) = 1$.

We have $\text{rank}(E(K)) = 0$, so

$$\text{rank}(E^L(L)) = \text{rank}(E(K)) + \text{rank}(E^L(K)) = \text{rank}(E^L(K)),$$

and if Conjecture III $T_2(K)$ holds then $\text{rank}(E^L(K)) = 1$. \square

7. TWO-DESCENTS OVER CYCLIC EXTENSIONS OF ODD PRIME DEGREE

Fix for this section a number field K , and a cyclic extension L/K of prime degree $p > 2$. Let $G = \text{Gal}(L/K)$. If R is a commutative ring, let $R[G]^0$ denote the augmentation ideal in the group ring $R[G]$.

Since $|G|$ is odd, the group ring $\mathbf{F}_2[G]$ is an étale \mathbf{F}_2 -algebra. Concretely, if we fix a generator of G we have G -isomorphisms

$$(12) \quad \mathbf{F}_2[G] \cong \mathbf{F}_2[X]/(X^p - 1) \cong \mathbf{F}_2 \oplus (\prod_{\pi} \mathbf{F}_2[X]/\pi(X))$$

where π runs through the irreducible factors of $X^{p-1} + \dots + 1$ in $\mathbf{F}_2[X]$, and the chosen generator of G acts on $\mathbf{F}_2[X]$ as multiplication by X . The submodule of $\mathbf{F}_2[G]$ corresponding to the summand \mathbf{F}_2 in (12) is $\mathbf{F}_2[G]^G$, and the submodule of $\mathbf{F}_2[G]$ corresponding to $\prod_{\pi} \mathbf{F}_2[X]/\pi(X)$ is the augmentation ideal $\mathbf{F}_2[G]^0$. Thus (12) corresponds to the decomposition (independent of choice of generator of G)

$$\mathbf{F}_2[G] = \mathbf{F}_2[G]^G \oplus \mathbf{F}_2[G]^0 = \mathbf{F}_2 \oplus (\oplus_{\mathbb{k} \in \Omega} \mathbb{k})$$

where Ω is the set of simple submodules of $\mathbf{F}_2[G]$ on which G acts nontrivially.

If B is an $\mathbf{F}_2[G]$ -module, then $B \otimes_{\mathbf{F}_2[G]} \mathbf{F}_2 = B^G$, and we define

$$B^{\text{new}} = B \otimes_{\mathbf{F}_2[G]} \mathbf{F}_2[G]^0 = \oplus_{\mathbb{k} \in \Omega} (B \otimes_{\mathbf{F}_2[G]} \mathbb{k}).$$

This gives a canonical decomposition $B = B^G \oplus B^{\text{new}}$.

Suppose now that E is an elliptic curve over K . The 2-Selmer group $\text{Sel}_2(E/L)$ has a natural action of $\mathbf{F}_2[G]$. Since $|G|$ is odd, it is straightforward to check that $\text{Sel}_2(E/L)^G = \text{Sel}_2(E/K)$, so

$$\text{Sel}_2(E/L) = \text{Sel}_2(E/K) \oplus \text{Sel}_2(E/L)^{\text{new}}.$$

Further, for $\mathbb{k} \in \Omega$ we define

$$\text{Sel}_{\mathbb{k}}(E/L) := \text{Sel}_2(E/L) \otimes_{\mathbf{F}_2[G]} \mathbb{k}, \quad d_{\mathbb{k}}(E/L) := \dim_{\mathbb{k}} \text{Sel}_{\mathbb{k}}(E/L)$$

so $\text{Sel}_2(E/L)^{\text{new}} = \oplus_{\mathbb{k} \in \Omega} \text{Sel}_{\mathbb{k}}(E/L)$.

Remark. Our proof of Theorem 1.11 for L/K goes as follows. We show that if E satisfies the hypotheses of Theorem 1.4, then:

- (1) There is a twist E' of E over K such that $d_{\mathbb{k}}(E'/L) = 0$ for some \mathbb{k} (see Proposition 7.3).
- (2) For every $r \geq 0$, there is a twist E' of E over K such that $d_2(E'/K) = r$ and $\text{Sel}_2(E'/L)^{\text{new}} = \text{Sel}_2(E/L)^{\text{new}}$ (see Proposition 7.4). In other words, we can twist to get whatever size we want for the “old part” of Selmer, while keeping the “new part” of Selmer unchanged.

Replacing E by a quadratic twist as necessary, by (1) we may assume $d_{\mathbb{k}}(E/L) = 0$ for some \mathbb{k} . Then by (2) we may assume that both $d_2(E/K) = 1$ and $d_{\mathbb{k}}(E/L) = 0$. Since $d_{\mathbb{k}}(E/L) = 0$ for some \mathbb{k} , we have $\text{rank}(E(L)) = \text{rank}(E(K))$ (see Lemma 7.1), and if Conjecture III $T_2(K)$ holds, then $\text{rank}(E(K)) = 1$.

Lemma 7.1. *Suppose E is an elliptic curve over K . If $d_{\mathbb{k}}(E/L) = 0$ for some $\mathbb{k} \in \Omega$, then $\text{rank}(E(L)) = \text{rank}(E(K))$.*

Proof. Since G is cyclic of prime order, it has only 2 irreducible rational representations, namely \mathbf{Q} (the trivial representation) and the augmentation ideal $\mathbf{Q}[G]^0$. Therefore we have an isomorphism of G -modules

$$E(L) \otimes \mathbf{Q} \cong \mathbf{Q}^a \times (\mathbf{Q}[G]^0)^b$$

for some $a, b \geq 0$. Then $E(L)$ has a submodule isomorphic to $(\mathbf{Z}[G]^0)^b$, so $E(L) \otimes \mathbf{Z}_2$ has a direct summand isomorphic to $(\mathbf{Z}_2[G]^0)^b$, so $E(L) \otimes \mathbf{F}_2$ has a submodule isomorphic to $(\mathbf{F}_2[G]^0)^b$, which implies that $d_{\mathbb{k}}(E/L) \geq b$. Since $d_{\mathbb{k}}(E/L) = 0$ we have $b = 0$, and so $\text{rank}(E(L)) = \text{rank}(E(K)) = a$. \square

We will need the following G -equivariant version of Proposition 3.3.

Proposition 7.2. *Suppose F/K is a quadratic extension and the hypotheses of Proposition 3.3 are satisfied. Let T be the set of primes of K where F/K is ramified, and T_L is the set of primes of L above T .*

(i) *If the localization map $\text{loc}_{T_L} : \text{Sel}_2(E/L)^{\text{new}} \rightarrow (\oplus_{\mathfrak{p} \in T_L} H_f^1(L_{\mathfrak{p}}, E[2]))^{\text{new}}$ is surjective, then there is an exact sequence*

$$0 \longrightarrow \text{Sel}_2(E^F/L)^{\text{new}} \longrightarrow \text{Sel}_2(E/L)^{\text{new}} \xrightarrow{\text{loc}_{T_L}} (\oplus_{\mathfrak{p} \in T_L} H_f^1(L_{\mathfrak{p}}, E[2]))^{\text{new}} \longrightarrow 0.$$

(ii) *Suppose that for every prime $\mathfrak{p} \in T$, \mathfrak{p} is inert in L/K and $E(K_{\mathfrak{p}})[2] \neq 0$. Then $\text{Sel}_2(E^F/L)^{\text{new}} = \text{Sel}_2(E/L)^{\text{new}}$.*

Proof. The proof is identical to that of Proposition 3.3, using that the functor $B \mapsto B^{\text{new}}$ is exact on $\mathbf{F}_2[G]$ -modules. As in the proof of Proposition 3.3, we have (G -equivariant) exact sequences

$$(13) \quad 0 \longrightarrow \mathcal{S}_{T_L}^{\text{new}} \longrightarrow \text{Sel}_2(E/L)^{\text{new}} \xrightarrow{\text{loc}_{T_L}} (\oplus_{\mathfrak{p} \in T_L} H_f^1(L_{\mathfrak{p}}, E[2]))^{\text{new}}$$

$$(14) \quad 0 \longrightarrow \mathcal{S}_{T_L}^{\text{new}} \longrightarrow \text{Sel}_2(E^F/L)^{\text{new}} \longrightarrow (\oplus_{\mathfrak{p} \in T_L} H_f^1(L_{\mathfrak{p}}, E^F[2]))^{\text{new}}$$

either of which can be taken as the definition of $\mathcal{S}_{T_L}^{\text{new}}$. The proof of Proposition 3.3 showed that if loc_{T_L} is surjective, then the right-hand map of (14) is zero, and then (13) is the exact sequence of (i).

Suppose $\mathfrak{p} \in T$ is inert in L/K . Let $\text{Frob}_{\mathfrak{p}} \in \text{Gal}(K_{\mathfrak{p}}^{\text{ur}}/K_{\mathfrak{p}})$ be the Frobenius of \mathfrak{p} , so $\text{Frob}_{\mathfrak{p}}^p = \text{Frob}_{\mathfrak{p}}^p$ is the Frobenius of the prime \mathfrak{P} above \mathfrak{p} . Since $\mathfrak{p} \in T$, the hypotheses of Proposition 3.3 require that E has good reduction at \mathfrak{p} , so by (1) there is a commutative diagram with horizontal isomorphisms

$$(15) \quad \begin{array}{ccc} H_f^1(L_{\mathfrak{P}}, E[2]) & \xrightarrow{\sim} & E[2]/(\text{Frob}_{\mathfrak{P}} - 1)E[2] \\ \text{Res} \uparrow & & \uparrow \\ H_f^1(K_{\mathfrak{p}}, E[2]) & \xrightarrow{\sim} & E[2]/(\text{Frob}_{\mathfrak{p}} - 1)E[2]. \end{array}$$

If $E(K_{\mathfrak{p}})[2] \neq 0$, then $\text{Frob}_{\mathfrak{p}}$ acts on $E[2]$ as an element of order 1 or 2, so $\text{Frob}_{\mathfrak{P}}|_{E[2]} = \text{Frob}_{\mathfrak{p}}|_{E[2]}$ and the groups on the right have the same order. The left-hand vertical map is injective since $[L_{\mathfrak{P}} : K_{\mathfrak{p}}]$ is odd. Therefore the left-hand map is an isomorphism, so G acts trivially on $H_f^1(L_{\mathfrak{P}}, E[2])$, and we have $H^1(L_{\mathfrak{P}}, E[2])^{\text{new}} = 0$.

If every $\mathfrak{p} \in T$ has these properties, then $(\oplus_{\mathfrak{p} \in T_L} H_f^1(L_{\mathfrak{p}}, E[2]))^{\text{new}} = 0$, so (ii) follows from (i). \square

Proposition 7.3. *Suppose E is an elliptic curve over K satisfying the hypotheses of Theorem 1.4. If $\text{Sel}_{\mathbb{k}}(E/L) \neq 0$ for every $\mathbb{k} \in \Omega$, then there is a quadratic twist E' of E over K such that*

$$d_{\mathbb{k}}(E'/L) = d_{\mathbb{k}}(E/L) - 1$$

for every $\mathbb{k} \in \Omega$.

Proof. Let Δ be the discriminant of (some integral model of) E . Let $M := K(E[2])$, so M is an S_3 -extension of K containing the quadratic extension $K(\sqrt{\Delta})$. Let Σ be the set of all infinite places and all primes where E has bad reduction.

Let \mathfrak{d} be the (formal) product of all places in $\Sigma - \{v_0\}$, where $v_0 \nmid 2$ is the distinguished place of Theorem 1.4, either real with $\Delta_{v_0} < 0$, or of multiplicative reduction with $\text{ord}_{v_0}(\Delta)$ odd. Let $K(8\mathfrak{d})$ denote the ray class field of K modulo $8\mathfrak{d}$, and let $K[8\mathfrak{d}]$ denote the maximal 2-power extension of K in $K(8\mathfrak{d})$. Note that $K(\sqrt{\Delta})/K$ is ramified at v_0 , but $LK[8\mathfrak{d}]/K$ is unramified at v_0 , so $M \cap LK[8\mathfrak{d}] = K(\sqrt{\Delta}) \cap LK[8\mathfrak{d}] = K$. Fix an element $\sigma \in G_K$, trivial on $LK[8\mathfrak{d}]$, whose projection to $\text{Gal}(MLK[8\mathfrak{d}]/LK[8\mathfrak{d}]) \cong \text{Gal}(M/K) \cong S_3$ has order 2. Since σ has order 2 on M , we have $E[2]/(\sigma - 1)E[2] \cong \mathbf{Z}/2\mathbf{Z}$.

Since $\text{Sel}_{\mathbb{k}}(E/L) \neq 0$ for every $\mathbb{k} \in \Omega$, it follows that $\text{Sel}_2(E/L)^{\text{new}}$ has a submodule free of rank one over $\mathbf{F}_2[G]^0$. Let $C \subset \text{Sel}_2(E/L)^{\text{new}}$ be such a submodule, fix an isomorphism $\eta : C \rightarrow \mathbf{F}_2[G]^0$, and define $\phi : C \rightarrow E[2]/(\sigma - 1)E[2]$ by $\phi(c) = f_1(\eta(c))x$, where $f_1 : \mathbf{F}_2[G] \rightarrow \mathbf{F}_2$ is projection onto the first coefficient, i.e., $f_1(\sum a_g g) = a_1$, and x is the nonzero element of $E[2]/(\sigma - 1)E[2]$.

By Lemma 3.5 we can find $\gamma \in G_K$ such that

- $\gamma|_{LMK[8\mathfrak{d}]} = \sigma$,
- $c(\gamma) = \phi(c)$ for all $c \in C$.

Let N be a Galois extension of K containing $MLK[8\mathfrak{d}]$, large enough so that the restriction of c to N is zero. Let \mathfrak{p} be a prime of K where E has good reduction, not dividing 2, unramified in L/K , whose Frobenius in $\text{Gal}(N/K)$ is in the conjugacy class of γ . Since $\gamma|_{K[8\mathfrak{d}]} = \sigma|_{K[8\mathfrak{d}]} = 1$, and $[K(8\mathfrak{d}) : K[8\mathfrak{d}]]$ is odd, there is an odd positive integer h such that $\gamma^h|_{K(8\mathfrak{d})} = 1$. Therefore \mathfrak{p}^h is principal, with a generator $\pi \equiv 1 \pmod{8\mathfrak{d}}$, positive at all real embeddings different from v_0 . Let $F = K(\sqrt{\pi})$. Then all places v dividing 2 and all places in $\Sigma - \{v_0\}$ split in F , and F/K is ramified only at \mathfrak{p} . Let E' be the quadratic twist of E by F . We will show that E' has the desired properties.

We will apply Proposition 7.2. Let $T = \{\mathfrak{p}\}$, and T_L the set of primes of L above \mathfrak{p} . For every $\mathfrak{P} \in T_L$,

$$H_{\mathfrak{f}}^1(L_{\mathfrak{P}}, E[2]) = H^1(L_{\mathfrak{P}}^{\text{ur}}/L_{\mathfrak{P}}, E[2]) = E[2]/(\text{Frob}_{\mathfrak{P}} - 1)E[2] = E[2]/(\sigma - 1)E[2]$$

is a one-dimensional \mathbf{F}_2 -vector space. Fix a prime of N above \mathfrak{p} whose Frobenius in $\text{Gal}(N/K)$ is equal to γ , and let \mathfrak{P}_0 be the corresponding prime of L . Then $T_L = \{\mathfrak{P}_0^\tau : \tau \in G\}$, and $\text{Frob}_{\mathfrak{P}_0^\tau/\mathfrak{p}}(N/K) = \tau\gamma\tau^{-1}$. The localization map

$$\text{loc}_{T_L} : \text{Sel}_2(E/L) \rightarrow \bigoplus_{\mathfrak{P} \in T_L} H_{\mathfrak{f}}^1(L_{\mathfrak{P}}, E[2]) \cong \mathbf{F}_2[G] \otimes_{\mathbf{Z}} (E[2]/(\sigma - 1)E[2])$$

is given on $c \in C \subset \text{Sel}_2(E/L)^{\text{new}}$ by

$$\begin{aligned} \text{loc}_{T_L}(c) &= \sum_{\tau} \tau \otimes c(\tau\gamma\tau^{-1}) = \sum_{\tau} \tau \otimes c^{\tau^{-1}}(\gamma) = \sum_{\tau} \tau \otimes \phi(c^{\tau^{-1}}) \\ &= \sum_{\tau} \tau \otimes f_1(\tau^{-1}\eta(c))x = \sum_{\tau} \tau \otimes f_{\tau}(\eta(c))x = \eta(c) \otimes x \end{aligned}$$

where $f_{\tau} : \mathbf{F}_2[G] \rightarrow \mathbf{F}_2$ is the map $f_{\tau}(\sum a_g g) = a_{\tau}$. Since the image of η is $\mathbf{F}_2[G]^0$, this shows that the localization map $C \rightarrow (\bigoplus_{\mathfrak{P} \in T_L} H_{\mathfrak{f}}^1(L_{\mathfrak{P}}, E[2]))^{\text{new}}$ is surjective. Now the proposition follows from Proposition 7.2(i). \square

Proposition 7.4. *Suppose E is an elliptic curve over K satisfying the hypotheses of Theorem 1.4. Then:*

- (i) *There is a quadratic twist E' of E/K such that $d_2(E'/K) = d_2(E/K) + 1$ and $\text{Sel}_2(E'/L)^{\text{new}} = \text{Sel}_2(E/L)^{\text{new}}$.*
- (ii) *If $\text{Sel}_2(E/K) \neq 0$, then there is a quadratic twist E' of E/K such that $d_2(E'/K) = d_2(E/K) - 1$ and $\text{Sel}_2(E'/L)^{\text{new}} = \text{Sel}_2(E/L)^{\text{new}}$.*

Proof. Let Σ be the set of all places $v \mid 2\infty$ of K and all v of bad reduction, and let v_0 be the distinguished place of Theorem 1.4, either real with $\Delta_{v_0} < 0$, or of multiplicative reduction with $\text{ord}_{v_0}(\Delta)$ odd. By Proposition 5.1, for (i) or (ii) we can find a quadratic extension F/K satisfying

- $d_2(E^F/K) = d_2(E/K) + 1$ in (i), $d_2(E^F/K) = d_2(E/K) - 1$ in (ii),
- all $v \in \Sigma - \{v_0\}$ split in F/K , and v_0 is unramified in F/K ,
- F/K is ramified at exactly one prime \mathfrak{p} , $\mathfrak{p} \nmid 2$, \mathfrak{p} is inert in L/K , and $E(K_{\mathfrak{p}})[2] \cong \mathbf{Z}/2\mathbf{Z}$.

By Proposition 7.2(ii) applied with $T = \{\mathfrak{p}\}$, $\text{Sel}_2(E^F/L)^{\text{new}} = \text{Sel}_2(E/L)^{\text{new}}$ in both cases. \square

Corollary 7.5. *Suppose E/K satisfies the hypotheses of Theorem 1.4, and $r \geq 0$. Then there is a twist E' of E such that*

$$d_2(E'/K) = r, \quad \text{rank}(E'(L)) = \text{rank}(E'(K)).$$

Proof. Using Proposition 7.3 repeatedly, we can find a twist E'' of E such that $d_{\mathbb{k}}(E''/L) = 0$ for at least one \mathbb{k} . Then applying Proposition 7.4 repeatedly, we can find another twist E' of E such that $d_{\mathbb{k}}(E'/L) = 0$ and $d_2(E'/K) = r$. Now the corollary follows from Lemma 7.1. \square

Proof of Theorem 1.11. Let $p = [L : K]$. If $p = 2$, Theorem 1.11 is Theorem 6.2, so we may assume that p is odd. By Lemma 5.4, we can find an elliptic curve E over K and a prime $\mathfrak{p} \nmid 2$ such that E has multiplicative reduction at \mathfrak{p} , $\text{ord}_{\mathfrak{p}}(\Delta_E) = 1$, and $\text{Gal}(K(E[2])/K) \cong S_3$. Then E satisfies the hypotheses of Theorem 1.4, so by Corollary 7.5, E has a twist with the desired properties. \square

8. PROOF OF THEOREM 1.12

The proof of Theorem 1.12 from Theorem 1.11 is due to Bjorn Poonen and Alexandra Shlapentokh. We thank them for allowing us to include their ideas here.

Definition 8.1. Suppose that R is a commutative ring with identity. Following [DL, De], we say that a subset D of R is *diophantine over R* if there is a finite set of polynomials $f_1, \dots, f_k \in R[X, Y_1, \dots, Y_m]$ for some m such that for every $x \in R$,

$$x \in D \iff \text{for } 1 \leq i \leq k \text{ there are } y_{1,i}, \dots, y_{m,i} \in R \\ \text{such that } f_i(x, y_{1,i}, \dots, y_{m,i}) = 0 \text{ for } 1 \leq i \leq k.$$

Lemma 8.2 ([DL]). *Suppose $K \subset L$ are number fields. Then:*

- (i) *If $D_1, D_2 \subset \mathcal{O}_L$ are diophantine over \mathcal{O}_L , then so is $D_1 \cap D_2$.*
- (ii) *If $D \subset \mathcal{O}_K$ is diophantine over \mathcal{O}_K , and \mathcal{O}_K is diophantine over \mathcal{O}_L , then D is diophantine over \mathcal{O}_L .*
- (iii) *If \mathbf{Z} is diophantine over \mathcal{O}_L , then \mathbf{Z} is diophantine over \mathcal{O}_K .*

Proof. This is Proposition 1(a), (c), and (d) of [DL]. \square

Corollary 8.3. *Suppose L/K is a cyclic extension of number fields. If Conjecture $\text{III}T_2(F)$ holds for all subfields $F \subset L$, then \mathcal{O}_K is diophantine over \mathcal{O}_L .*

Proof. We have $K = K_0 \subset K_1 \subset \cdots \subset K_n = L$, where each K_{i+1}/K_i is cyclic of prime degree. If Conjecture $\text{III}T_2(K_i)$ holds for every i , then by Theorem 1.11 for every i there is an elliptic curve E/K_i such that $\text{rank}(E(K_i)) = \text{rank}(E(K_{i+1})) = 1$. By Theorem 1 of [Po], it follows that \mathcal{O}_{K_i} is diophantine over $\mathcal{O}_{K_{i+1}}$. Now the corollary follows from Lemma 8.2(ii) by induction. \square

Proof of Theorem 1.12. Fix a number field K , and let L be the Galois closure of K/\mathbf{Q} . For every $g \in \text{Gal}(L/\mathbf{Q})$, let $L^{(g)}$ denote the fixed field of g in L . Then $L/L^{(g)}$ is cyclic, so $\mathcal{O}_{L^{(g)}}$ is diophantine over \mathcal{O}_L by Corollary 8.3. But then by Lemma 8.2(i), $\cap_g \mathcal{O}_{L^{(g)}} = \mathcal{O}_L^{\text{Gal}(L/\mathbf{Q})} = \mathbf{Z}$ is diophantine over \mathcal{O}_L , so by Lemma 8.2(iii), \mathbf{Z} is diophantine over \mathcal{O}_K . Now the theorem follows from Matiyasevich's Theorem [Mat]. \square

9. ELLIPTIC CURVES WITH CONSTANT PARITY

In this section we discuss briefly the phenomenon of “constant parity”.

Definition 9.1. Suppose E is an elliptic curve defined over a number field K . We will say that E/K has *constant 2-Selmer parity* if the parity of $d_2(E^F/K)$ is constant as F ranges over all quadratic extensions of K , i.e., if $d_2(E^F/K) \equiv d_2(E/K) \pmod{2}$ for all quadratic extensions F/K .

Similarly, we can say that E has constant Mordell-Weil parity if the parity of $\text{rank}(E^F(K))$ is independent of the quadratic extension F/K , and E has constant analytic parity if the global root number of E^F/K is independent of F . Standard conjectures imply that all three notions of constant parity are the same.

Example 9.2. Suppose E has complex multiplication by an imaginary quadratic field $k \subset K$. Then E has constant (even) 2-Selmer parity, constant (even) Mordell-Weil parity, and constant (even) analytic parity.

The question of constant analytic parity was studied by T. Dokchitser and V. Dokchitser in [DD]. They proved the following.

Theorem 9.3 (Theorem 1 of [DD]). *An elliptic curve E over a number field K has constant analytic parity if and only if K is totally imaginary and E acquires good reduction over an abelian extension of K .*

The following example from [DD] shows that constant parity can be odd.

Example 9.4. Suppose K is totally imaginary, E/K has good reduction everywhere, and $[K : \mathbf{Q}]/2$ is odd. Then E/K has constant odd analytic parity (see [Ro, Theorem 2(i) and Proposition 8(i)]).

This applies in particular to the elliptic curve $E : y^2 + xy = x^3 + x^2 - 2x - 7$ (labelled 121C1 in Cremona's tables) and K the splitting field of $x^3 - 11$.

From now on we will only consider constant 2-Selmer parity. The following theorem will be proved at the end of this section.

Theorem 9.5. *If E/K has constant 2-Selmer parity, then K is totally imaginary and E has additive reduction at all primes.*

Definition 9.6. Suppose E is an elliptic curve defined over a local field K . If F is a quadratic extension of K (or $F = K$), define

$$\delta(E, F/K) = \dim_{\mathbf{F}_2} E(K)/\mathbf{N}_{F/K}E(F).$$

We will say that E/K has *constant local parity* if $\delta(E, F/K)$ is even for every quadratic extension F/K .

If $D \in K^\times/(K^\times)^2$, we will say that E/K has *D -parity* if

$$\delta(E, F/K) \text{ is even} \iff D \in \mathbf{N}_{F/K}F^\times.$$

Note that if D is a square in K^\times , then E/K has D -parity if and only if it has constant local parity.

Lemma 9.7. *Suppose E is an elliptic curve defined over a local field K , and $\Delta_E \in K^\times/(K^\times)^2$ is its discriminant.*

- (i) *If v is nonarchimedean with residue characteristic different from 2, and E has good reduction, then E has Δ_E -parity.*
- (ii) *If K is nonarchimedean and E has multiplicative reduction, then E does not have Δ_E -parity.*
- (iii) *If $K = \mathbf{R}$, then E does not have Δ_E -parity.*

Proof. Assertions (i), (ii), and (iii) are [Maz, Corollary 4.4] and [Kr, Proposition 3],[Kr, Propositions 1 and 2], and [Kr, Proposition 6], respectively. \square

For the rest of this section, fix an elliptic curve E defined over a number field K , and let Δ_E be the discriminant of some model of E .

- Theorem 9.8.**
- (i) *If E/K_v has constant local parity for every place v of K , then E/K has constant 2-Selmer parity.*
 - (ii) *E/K has constant 2-Selmer parity if and only if E/K_v has Δ_E -parity for every v .*

Proof. Suppose F is a quadratic extension of K . Kramer's congruence (Theorem 2.3) says

$$(16) \quad d_2(E^F/K) \equiv d_2(E/K) + \sum_v \delta(E, F_v/K_v) \pmod{2}$$

where F_v is the completion of F at some place above v . Assertion (i) follows directly from this.

Now suppose E/K_v has Δ_E -parity for every v . Then, if τ is the nontrivial automorphism of $\text{Gal}(F/K)$,

$$\tau^{\delta(E, F_v/K_v)} = [\Delta_E, F_v/K_v]$$

where $[\cdot, F_v/K_v]$ is the local Artin symbol. The global reciprocity law shows that $\prod_v [\Delta_E, F_v/K_v] = 1$, so $\sum_v \delta(E, F_v/K_v)$ is even and it follows from (16) that E/K has constant 2-Selmer parity.

Finally, suppose that for some v_0 , E/K_{v_0} does not have Δ_E -parity. By Lemma 9.7(i), E/K_v has Δ_E -parity for almost all v . Fix a quadratic extension F/K such that

- $\tau^{\delta(E, F_{v_0}/K_{v_0})} = \tau \cdot [\Delta_E, F_{v_0}/K_{v_0}]$,
- every $v \neq v_0$ where E/K_v does not have Δ_E -parity splits in F/K .

Then $\tau^{\delta(E, F_v/K_v)} = [\Delta_E, F_v/K_v]$ for every $v \neq v_0$, so

$$\tau^{\sum_v \delta(E, F_v/K_v)} = \tau \cdot \prod_v [\Delta_E, F_v/K_v] = \tau,$$

so by (16), $d_2(E/K)$ and $d_2(E^F/K)$ have opposite parity. \square

Proof of Theorem 9.5. Theorem 9.5 follows directly from Theorem 9.8(ii) and Lemma 9.7(ii,iii). \square

Corollary 9.9. *If Δ_E is a square, then E/K has constant 2-Selmer parity if and only if E/K_v has constant local parity for every v .*

Proof. This is immediate from Theorem 9.8(ii). \square

REFERENCES

- [Ca1] J.W.S. Cassels, Arithmetic on an elliptic curve. Proc. Internat. Congr. Mathematicians (Stockholm, 1962) Djursholm: Inst. Mittag-Leffler (1963) 234–246.
- [Ca2] J.W.S. Cassels, Arithmetic on curves of genus 1. VIII. On conjectures of Birch and Swinnerton-Dyer. *J. Reine Angew. Math.* **217** (1965) 180–199.
- [Ch1] S. Chang, On the arithmetic of twists of superelliptic curves. *Acta Arith.* **124** (2006) 371–389.
- [Ch2] S. Chang, Quadratic twists of elliptic curves with small Selmer rank. Preprint available at <http://arxiv.org/abs/0809.5019>.
- [De] J. Denef, Diophantine sets over algebraic integer rings. II. *Trans. Amer. Math. Soc.* **257** (1980) 227–236.
- [DL] J. Denef, L. Lipshitz, Diophantine sets over some rings of algebraic integers. *J. London Math. Soc.* **18** (1978) 385–391.
- [DD] T. Dokchitser, V. Dokchitser, Elliptic curves with all quadratic twists of positive rank. *Acta Arith.* **137** (2009) 193–197.
- [GM] F. Gouvêa, B. Mazur, The square-free sieve and the rank of elliptic curves. *J. Amer. Math. Soc.* **4** (1991) 1–23.
- [HB] D.R. Heath-Brown, The size of Selmer groups for the congruent number problem II. *Invent. Math.* **118** (1994) 331–370.
- [Kr] K. Kramer, Arithmetic of elliptic curves upon quadratic extension, *Transactions Amer. Math. Soc.* **264** (1981) 121–135.
- [Mat] Yu. V. Matiyasevich, The Diophantineness of enumerable sets. *Dokl. Akad. Nauk SSSR* **191** (1970) 279–282.
- [Maz] B. Mazur, Rational points of abelian varieties with values in towers of number fields. *Invent. Math.* **18** (1972) 183–266.
- [MR1] B. Mazur and K. Rubin, Kolyvagin systems. *Memoirs of the Amer. Math. Soc.* **799** (2004).
- [MR2] B. Mazur, K. Rubin, Finding large Selmer rank via an arithmetic theory of local constants. *Annals of Math.* **166** (2007) 581–614.
- [Me] L. Merel, Bornes pour la torsion des courbes elliptiques sur les corps de nombres. *Invent. Math.* **124** (1996) 437–449.
- [Mi] J.S. Milne, Arithmetic duality theorems, *Perspectives in Math.* **1**, Orlando: Academic Press (1986).
- [On] K. Ono, Nonvanishing of quadratic twists of modular L -functions and applications to elliptic curves. *J. Reine Angew. Math.* **533** (2001) 81–97.
- [OS] K. Ono, C. Skinner, Non-vanishing of quadratic twists of modular L -functions. *Invent. Math.* **134** (1998) 651–660.
- [Po] B. Poonen, Using elliptic curves of rank one towards the undecidability of Hilbert’s tenth problem over rings of algebraic integers. In: Algorithmic Number Theory (Sydney, 2002), *Lecture Notes in Comput. Sci.* **2369**, Berlin: Springer-Verlag (2002) 33–42.
- [Ro] D. Rohrlich, Galois theory, elliptic curves, and root numbers, *Compositio Math.* **100** (1996) 311–349.
- [Ru] K. Rubin, Euler Systems. *Annals of Math. Studies* **147**, Princeton: Princeton University Press (2000).

- [Se] J-P. Serre, Divisibilité de certaines fonctions arithmétiques. *Enseignement Math.* **22** (1976) 227–260.
- [Sh] A. Shlapentokh, Hilbert’s tenth problem over number fields, a survey. In: Hilbert’s tenth problem: relations with arithmetic and algebraic geometry (Ghent, 1999), *Contemp. Math.* **270**, Amer. Math. Soc.: Providence (2000) 107–137.
- [S1] J.H. Silverman, The Arithmetic of Elliptic Curves, Graduate Texts in Mathematics **106**, New York: Springer-Verlag (1986).
- [S2] J.H. Silverman, Advanced Topics in the Arithmetic of Elliptic Curves, Graduate Texts in Mathematics **151**, New York: Springer-Verlag (1994).
- [SD] H.P.F. Swinnerton-Dyer, The effect of twisting on the 2-Selmer group. *Math. Proc. Cambridge Philos. Soc.* **145** (2008) 513–526.
- [T1] J. Tate, Duality theorems in Galois cohomology over number fields, in: *Proc. Intern. Cong. Math.*, Stockholm (1962) 234–241.
- [T2] J. Tate, Algorithm for determining the type of a singular fiber in an elliptic pencil. In: Modular functions of one variable (IV), *Lecture Notes in Math.* **476**, New York: Springer-Verlag (1975) 33–52.
- [Wi] A. Wintner, On the prime number theorem. *Amer. J. Math.* **64** (1942) 320–326.

DEPARTMENT OF MATHEMATICS, HARVARD UNIVERSITY, CAMBRIDGE, MA 02138, USA
E-mail address: `mazur@math.harvard.edu`

DEPARTMENT OF MATHEMATICS, UC IRVINE, IRVINE, CA 92697, USA
E-mail address: `krubin@math.uci.edu`