

# Energy and length in a topological planar quadrilateral

Sa'ar Hersensky \*

## Abstract

We provide bounds for the product of the lengths of distinguished shortest paths in a finite network induced by a triangulation of a topological planar quadrilateral.

## 1 Introduction

A topological planar closed disk with four distinguished points on its boundary, its corners, will be called a quadrilateral. The following definition is due to Schramm ([16]).

**Definition 1.1** Let  $Q$  be a quadrilateral endowed with a triangulation. Let  $V, E, T$  denote the set of vertices, edges and triangles of  $Q$ , respectively. Let  $\partial Q = P_1 \cup P_2 \cup P_3 \cup P_4$  be a decomposition of  $\partial Q$  into four non-trivial arcs of the triangulation with disjoint interiors, in cyclic order. If the intersection of two any of these arcs is not empty, then it consists of a corner (all of which are vertices). A corner may belong to one and only one of the  $P_i$ 's. The collection  $\mathcal{T} = (V, E, T, P_1, P_2, P_3, P_4)$  will be called a triangulation of  $Q$ .

By invoking a *conductance function*,  $\mathcal{T}$  becomes a *finite network*. One can define a *boundary value problem* (BVP) on the network. Let  $f$  be the solution of the BVP and let  $I(f)$  be its *Dirichlet energy*. Corollary 3.7 provides inequalities relating the product of the lengths of a shortest *thick* vertical path (a path which connects  $P_3$  and  $P_1$ , see Definition 3.4) and a shortest thick horizontal path (a path which connects  $P_2$  and  $P_4$ , see Definition 3.4) in terms of  $I(f)$  and some constants arising from the combinatorics and the conductance function. Corollary 3.7 follows from Theorem 3.5 (our main theorem) and Lemma 3.6. The length is measured with respect to  $\rho$ , the *gradient metric* (Definition 3.3) which is induced by the solution of the BVP (see Section 2 and Section 3 for the details). In the special case where  $c(x, y) \equiv 1$  and  $k$  is the maximal degree of  $x \in V$ , it follows from Corollary 3.7 that

$$l(|V|)I(f) \geq \text{length}_\rho(\gamma^*)\text{length}_\rho(\gamma) \geq \frac{1}{\sqrt{k}}I(f), \text{ where } l(|V|) \text{ is some constant.}$$

C. Loewner (see [7]) studied Differential-geometric inequalities relating area and the product of shortest (vertical and horizontal) curves in a quadrilateral. His inequalities are derived with respect to the Euclidean metric. His work was generalized and forms a rich theory. The well known reciprocal property of the *extremal lengths* of conjugate families

---

\*The research is partially supported by The David & Elaine Potter Charitable Foundation

of curves in a quadrilateral is one useful example of this theory (see [1], [14] and [15] for a few examples and generalizations for other Riemann surfaces). More recently, Cannon introduced a notion of extremal length in a discrete setting ([8]). His definition is distinct from the original notion of extremal length that was introduced by Duffin ([11]).

In the work of Cannon, Floyd and Parry (see for example [9]), as well as in the work of Schramm ([16]), inequalities generalizing the reciprocal property of the extremal length (with respect to some extremal metric) of conjugate families of curves in a quadrilateral are very useful.

In the setting of finite and infinite networks reciprocal property of extremal length and *capacity* were studied extensively with respect to an extremal metric (see [18] for a detailed account).

One motivation for using the gradient metric in this paper arises from extremal length arguments in the complex plane. It is well known (see [1]) that for every  $z \in \mathbf{C}$  the extremal metric in a topological quadrilateral in the complex plane satisfies

$$m(z) = |\nabla(f)(z)|,$$

where  $f$  is the solution of the classical Dirichlet-Neumann boundary value problem. In the complex plane it is also known that equality holds in Corollary 3.7, where both sides equal  $I(f)$ .

Consider the broader class of BVP problems that are studied in [2]. Our work is also motivated by the following.

**Question.** Is there a BVP problem and a metric  $\rho_0$  (which is perhaps different than  $\rho$ ) derived from the solution such that

$$\text{length}_{\rho_0}(\gamma^*)\text{length}_{\rho_0}(\gamma) \geq I(f) \text{ and } \sum_{x \in \bar{F}} \rho_0^2(x) = I(f) \text{ (see Definition 3.2)?}$$

**Remark.** In paper [13] we will use some of the ideas of this paper to prove a finite Riemann mapping theorem ([9], [16]). A more direct proof would follow from a positive answer to the question above. A finite Riemann mapping theorem can be viewed as the first step in solving the Cannon conjecture: A negatively curved group  $G$  with  $\partial G = S^2$  is Kleinian. We hope that our ideas will be useful towards the resolution of this conjecture.

**Acknowledgement.**

Part of this research was done while the author visited the IAS, the Mathematics departments of Princeton and Caltech during the summer of 2003. We express our deepest gratitude to both for their generous hospitality. We are grateful to Francis Bonahon, Nikolai Makarov and Steven Kerckhoff for enjoyable and helpful discussions. We are indebted to David Gabai and Frédéric Paulin for their careful reading of this work and many useful remarks. We are indebted to anonymous referees of this paper. Their remarks and corrections were essential.

## 2 Preliminaries

We recall some known facts regarding harmonic functions and boundary value problems on networks. We use the notations of Section 2 in [2]. Let  $\Gamma = (V, E, c)$  be a *finite network*, that is a simple and finite connected graph with a vertex set  $V$  and edge set

$E$ . We shall also assume that the graph is planar. Each edge  $(x, y) \in E$  is assigned a *conductance*  $c(x, y) = c(y, x) > 0$ . Let  $\mathcal{P}(V)$  denote the set of non-negative functions on  $V$ . If  $u \in \mathcal{P}(V)$ , its support is given by  $S(u) = \{x \in V : u(x) \neq 0\}$ . Given  $F \subset V$  we denote by  $F^c$  its complement in  $V$ . Set  $\mathcal{P}(F) = \{u \in \mathcal{P}(V) : S(u) \subset F\}$ . The set  $\partial F = \{(x, y) \in E : x \in F, y \in F^c\}$  is called the *edge boundary* of  $F$  and the set  $\delta F = \{x \in F^c : (x, y) \in E \text{ for some } y \in F\}$  is called the *vertex boundary* of  $F$ . Let  $\bar{F} = F \cup \delta F$  and let  $\bar{E} = \{(x, y) \in E : x \in F\}$ . Given  $F \subset V$ , let  $\bar{\Gamma}(F) = (\bar{F}, \bar{E}, \bar{c})$  be the network such that  $\bar{c}$  is the restriction of  $c$  to  $\bar{E}$ . We say that  $x \sim y$  if  $(x, y) \in \bar{E}$ . For  $x \in \bar{F}$  let  $k(x)$  denote the degree of  $x$  (if  $x \in \delta(F)$  the neighbors of  $x$  are taken only from  $F$ ).

For  $f, h : \bar{E} \rightarrow \mathbf{R}$  we let  $(f, h) = \sum_{(x,y) \in \bar{E}} \frac{f(x,y)h(x,y)}{c(x,y)}$  be an inner product on  $l^2(\bar{E}, 1/c)$  (see [19, 1.2.A]). The following definitions are discrete analogues of classical notions in continuous Potential Theory [12].

**Definition 2.1** ([3, Section 3]) Let  $u \in \mathcal{P}(\bar{F})$ ,

1. then for  $x \in \bar{F}$ , the function  $\Delta u(x) = \sum_{y \sim x} c(x, y)(u(x) - u(y))$  is called the potential of  $u$  at  $x$ , (if  $x \in \delta(F)$  the neighbors of  $x$  are taken only from  $F$ ) and
2. the number  $I(u) = \sum_{x \in \bar{F}} \Delta u(x)u(x) = \sum_{(x,y) \in \bar{E}} c(x, y)(u(x) - u(y))^2$ , is called the *Dirichlet energy* of  $u$ .
3. A function  $u \in \mathcal{P}(\bar{F})$  is called harmonic in  $F \subset V$  if  $\Delta u(x) = 0$ , for all  $x \in F$ .

When  $c(x, y) \equiv 1$ , an easy computation shows that  $u$  is harmonic at a vertex  $x$  if and only if the value of  $u$  at  $x$  is the arithmetic average of the value of  $u$  on the neighbors of  $x$ .

When  $(x, y) \in \bar{E}$  let us denote by  $[x, y]$  the directed edge from  $x$  to  $y$  and let  $\vec{E} = \{[x, y] : (x, y) \in \bar{E}\}$  denote the set of all directed edges. Given  $u : V \rightarrow \mathbf{R}$  we define the *differential* or the *gradient* of  $u$  as  $du : \vec{E} \rightarrow \mathbf{R}$  by  $du[x, y] = c(x, y)(u(y) - u(x))$  for all  $[x, y] \in \vec{E}$  (see the notations of Section 2 in [6] for example). Note that if  $|\vec{E}| = m$ , then  $du$  can be identified with a vector in  $\mathbf{R}^m$ . It now follows by Definition 2.1 that for every function  $u : V \rightarrow \mathbf{R}$  we have that  $I(u) = \frac{1}{2} \sum_{e \in \vec{E}} du(e)^2$ .

let  $\vec{E}(x)$  denote the set of all edges of the form  $[x, y]$  which are in  $\vec{E}$ . Any  $g : \vec{E}(x) \rightarrow \mathbf{R}$  can be naturally viewed as an element in  $\mathbf{R}^{k(x)}$ . We will denote this vector space, with the restriction of the inner product on  $\vec{E}$ , by  $T_x$ . In particular we have

**Definition 2.2** Let  $u : \bar{F} \rightarrow \mathbf{R}$ . Let  $\vec{du}(x) = du|_{\vec{E}(x)}$ , denote the restriction of  $du$  to  $\vec{E}(x)$  (In particular,  $\vec{du}(x)$  can be viewed as a vector in  $T_x$ ).

For  $x \in \delta(F)$  let  $\{y_1, y_2, \dots, y_m\} \in F$  be its neighbors, enumerated in a cyclic order.

**Definition 2.3** The normal vector derivative at  $x \in \delta(F)$  is defined by

$\frac{\partial u}{\partial n_F}(x) = (c(x, y_1)(u(x) - u(y_1)), \dots, c(x, y_m)(u(x) - u(y_m)))$  and the conductance vector at  $x$  is defined by  $\vec{c}_{\delta(F)}(x) = (c(x, y_1), \dots, c(x, y_m))$ .

If  $x \in F$ ,  $\vec{c}_F(x)$  is defined similarly and the neighbors of  $x$  are taken in  $F \cup \delta(F)$ .

The following definition provides the discrete analogue of the continuous notion of *normal derivative*.

**Definition 2.4** ([10]) The normal derivative of  $u$  at a point  $x \in \delta F$  with respect to the set  $F$  is

$$\frac{\partial u}{\partial n_F}(x) = \sum_{y \sim x, y \in F} c(x, y)(u(x) - u(y)).$$

The following proposition establishes a discrete version of the first classical *Green identity*. It will be crucial in the proof of Theorem 3.5.

**Proposition 2.5** ([2, Prop. 3.1]) (**The first Green identity**) *Let  $F \subset V$  and  $u, v \in \mathcal{P}(\bar{F})$ . Then we have that*

$$\sum_{(x, y) \in \bar{E}} c(x, y)(u(x) - u(y))(v(x) - v(y)) = \sum_{x \in F} \Delta u(x)v(x) + \sum_{x \in \delta(F)} \frac{\partial u}{\partial n_F}(x)v(x).$$

**Remarks.**

1. In [2] a second Green identity is obtained. In this paper we will use only the one above.
2. In [5] (see in particular Section 2 and Section 3) a systematic study of discrete calculus on  $n$ -dimensional (uniform) grids of Euclidean  $n$ -space is provided. Their definition of a tangent space may be adopted to our setting and does not require the notion of directed edges. However, in [13] directed edges will play an important role.

### 3 Length estimates of shortest paths

Throughout this section  $\mathcal{T}$  will denote a fixed triangulation of a quadrilateral (see Definition 1.1). We will denote by  $F$  the set of vertices which do not belong to  $\partial Q$ . Hence,  $\delta(F)$  is the set of vertices that belong to  $P_1 \cup P_2 \cup P_3 \cup P_4$ . Let  $\{c(x, y)\}_{(x, y) \in \bar{E}}$  be a fixed conductance function and let  $\Gamma(\bar{F})$  be the associated network. We are interested in functions that solve a boundary value problem (BVP) on  $\Gamma(\bar{F})$ . The following definition is based on [2, Section 3] and [4, Section 4].

**Definition 3.1** Let  $g > 0$  be a constant. A Dirichlet-Neumann boundary value function is a function  $f \in \mathcal{P}(\bar{F})$  which satisfies the following:

1.  $f$  is harmonic in  $F$ ,
2.  $f|_{P_2} = 0$ ,
3.  $f|_{P_4} = g$ , for some constant  $g$ , and
4.  $\frac{\partial f}{\partial n_F}|_{P_1} = \frac{\partial f}{\partial n_F}|_{P_3} = 0$ .

**Remark.** The uniqueness and existence of a Dirichlet-Neumann boundary value function is provided by the nice and foundational work in [2, Section 3] and [4, Section 4]. In fact, their work provides a detailed framework for a broader class of boundary value problems on finite networks.

**Definition 3.2** ([8]) A metric on a finite network is a function  $\rho : V \rightarrow [0, \infty)$ .

In particular, the length of a path is given by integrating  $\rho$  along the path. When  $\rho \equiv 1$ , the familiar distance function on  $V \times V$  is obtained by setting  $\text{dist}(A, B) = \sum_{x \in \alpha} 1 - 1 = k$ , where  $\alpha = (x, x_1, \dots, x_k)$  is a path with the smallest possible number of vertices among all the paths connecting a vertex in  $A$  and a vertex in  $B$ . We now define the gradient metric which will be used in our estimates.

**Definition 3.3** Given  $f \in \mathcal{P}(\bar{F})$  the gradient metric induced by  $f \in \mathcal{P}(\bar{F})$  is defined by

$$\rho(x) = \begin{cases} \|\vec{df}(x)\| & \text{if } x \in F \\ \|\frac{\partial f}{\partial n_F}(x)\| & \text{if } x \in \delta(F). \end{cases}$$

Before turning to our main theorem, we will define the paths which are going to be considered. The definition below singles out two classes of paths. These classes are separated enough from  $\delta(F)$ .

**Definition 3.4** A path  $\beta = (x_0, x_1, \dots, x_n)$  will be called vertically (horizontally) thick if it satisfies the following:

1.  $x_0 = \beta \cap P_3$  ( $x_0 = \beta \cap P_2$ ) and  $x_n = \beta \cap P_1$  ( $x_n = \beta \cap P_4$ ) respectively,
2. neither  $x_0$  or  $x_n$  is a corner,
3. for all  $i = 1, \dots, n-1$ ,  $x_i \in F$ ,
4. for all  $i = 2, \dots, n-2$ ,  $\text{dist}(x_i, P_3 \cup P_4 \cup P_1) > 1$  (,
5.  $\text{dist}(x_1, P_1 \cup P_3 \cup P_4) = \text{dist}(x_{n-1}, P_1 \cup P_3 \cup P_4) \geq 1$  ( $\text{dist}(x_1, \delta(F)) = \text{dist}(x_{n-1}, \delta(F)) \geq 1$ ), and equality is attained uniquely for  $x_0$  and  $x_n$  respectively.

We now turn to our main theorem.

**Theorem 3.5** Let  $\mathcal{T}$  be a triangulation of a topological quadrilateral. Let  $\Gamma(\bar{F})$  be the associated network. Let  $f$  be the Dirichlet-Neumann boundary value function with  $g \neq 0$ . Let  $\rho$  be the gradient metric induced by  $f$ . Let

$$M = \max_{x \in F \cup \delta(F)} \{ \|\vec{c}_F(x)\|, \|\vec{c}_{\delta(F)}(x)\| \} \text{ and let } m = \min_{(x,y) \in \bar{E}} \sqrt{c(x,y)}.$$

1. If  $\gamma^*$  is a shortest vertical thick path which connects  $P_1$  to  $P_3$ , then

$$\text{length}_\rho(\gamma^*) \geq \frac{I(f)}{gM}, \text{ and}$$

2. if  $\gamma$  is a shortest horizontal thick path which connects  $P_2$  to  $P_4$ , then

$$\text{length}_\rho(\gamma) \geq gm.$$

**Proof.** Using properties (1)-(4) of  $f$  and the first Green identity (Proposition 2.5) with  $u = v = f$  we obtain that

$$(1) \quad I(f) = \sum_{x \in P_4} \frac{\partial f}{\partial n_F}(x) f(x).$$

Hence, by the definition of  $g$ , we have that

$$(2) \quad I(f) = g \left| \sum_{x \in P_4} \frac{\partial f}{\partial n_F}(x) \right|.$$

Let  $\gamma^*$  be a shortest thick path connecting  $P_3$  to  $P_1$ . It is clear that we may assume that  $\gamma^*$  is simple. Let  $\gamma^* \cap P_3 = x_0$  and  $\gamma^* \cap P_1 = x_n$ .




Figure 1 : A shortest thick vertical path between  $P_3$  and  $P_1$ .

Let  $V_{\gamma^*}$  denote the subset of  $F$  which is enclosed, in a cyclic order, by  $\gamma^*$ , a part of  $P_1$  (which we will denote by  $P_1(\gamma^*)$ ),  $P_4$  and part of  $P_3$  (which we will denote by  $P_3(\gamma^*)$ ). It follows from Definition 3.4 that  $V_{\gamma^*} \neq \emptyset$  and that  $\delta(V_{\gamma^*}) = \gamma^* \cup P_1(\gamma^*) \cup P_4 \cup P_3(\gamma^*)$ .

We now apply the first Green identity with  $u = f$  and the constant function  $v \equiv 1$  in  $V_{\gamma^*} \cup \delta(V_{\gamma^*})$ . We obtain that

$$(3) \quad \sum_{x \in \delta(V_{\gamma^*})} \frac{\partial f}{\partial n_{V_{\gamma^*}}}(x) = 0.$$

It follows by Definition 3.4 that for every  $x \in P_1(\gamma^*) \cup P_4 \cup P_3(\gamma^*)$  which is not in  $\gamma^*$  we have that

$$\frac{\partial f}{\partial n_{V_{\gamma^*}}}(x) = \frac{\partial f}{\partial n_F}(x).$$

By using the fourth property of Definition 3.1 and the triangle inequality we have that

$$(4) \quad \left| \sum_{x \in P_4} \frac{\partial f}{\partial n_{V_{\gamma^*}}}(x) \right| = \left| \sum_{x \in \gamma^*} \frac{\partial f}{\partial n_{V_{\gamma^*}}}(x) \right| \leq \sum_{x \in \gamma^*} \left| \frac{\partial f}{\partial n_{V_{\gamma^*}}}(x) \right|.$$

For every  $x \in \gamma^*$  (viewed as a vertex in  $\delta(V_{\gamma^*})$ ) we have that

$$\frac{\partial f}{\partial n_{V_{\gamma^*}}}(x) = (\vec{e}_{\delta(V_{\gamma^*})}(x), \overrightarrow{\frac{\partial f}{\partial n_{V_{\gamma^*}}}}(x)).$$

Hence by the Cauchy-Schwartz inequality we have that

$$(5) \quad \left| \frac{\partial f}{\partial n_{V_{\gamma^*}}}(x) \right| = \left| \langle \vec{c}_{\delta(V_{\gamma^*})}(x), \overrightarrow{\frac{\partial f}{\partial n_{V_{\gamma^*}}}}(x) \rangle \right| \leq \|\vec{c}_{\delta(V_{\gamma^*})}(x)\| \left\| \overrightarrow{\frac{\partial f}{\partial n_{V_{\gamma^*}}}}(x) \right\|.$$

It is also clear that for every  $x \in \gamma^*$  which is different from  $x_0$  or  $x_n$ , we have that

$$(6) \quad \|\vec{c}_{\delta(V_{\gamma^*})}(x)\| \leq \|\vec{c}_F(x)\| \quad \text{and} \quad \left\| \overrightarrow{\frac{\partial f}{\partial n_{V_{\gamma^*}}}}(x) \right\| \leq \rho(x).$$

If  $x = x_0$  or  $x = x_n$  we have that

$$(7) \quad \|\vec{c}_{\delta(V_{\gamma^*})}(x)\| \leq \|\vec{c}_{\delta(F)}(x)\| \quad \text{and} \quad \left\| \overrightarrow{\frac{\partial f}{\partial n_{V_{\gamma^*}}}}(x) \right\| \leq \rho(x).$$

Hence, by summing over all  $x \in \gamma^*$ , the definition of  $M$ , Equations (2), (4), (5), (6) and (7) we have that

$$(8) \quad \frac{I(f)}{gM} \leq \sum_{x \in \gamma^*} \rho(x) = \text{length}_\rho(\gamma^*),$$

which is first assertion of the theorem.

Let  $\gamma$  be a shortest thick path connecting  $P_2$  and  $P_4$ . It is clear that we may assume that  $\gamma$  is simple. Let  $\gamma \cap P_2 = x_0$  and let  $\gamma \cap P_4 = x_n$ .




Figure 2 : A shortest thick horizontal path between  $P_2$  and  $P_4$ .

By integrating  $\rho$  along  $\gamma$  we have that

$$\sum_{x \in \gamma} \rho(x) = \left\| \overrightarrow{\frac{\partial f}{\partial n_F}}(x_0) \right\| + \|df(x_1)\| + \|df(x_2)\| + \dots + \|df(x_{n-1})\| + \left\| \overrightarrow{\frac{\partial f}{\partial n_F}}(x_n) \right\|.$$

For each  $i = 1, \dots, n-1$  we have that

$$\|df(x_i)\| \geq \sqrt{c(x_i, x_{i+1})} |f(x_{i+1}) - f(x_i)|.$$

It is easy to see that  $\|\frac{\overrightarrow{\partial f}}{\partial n_F}(x_0)\| \geq \sqrt{c(x_0, x_1)}|f(x_1) - f(x_0)|$  and that  $\|\frac{\overrightarrow{\partial f}}{\partial n_F}(x_n)\| \geq \sqrt{c(x_{n-1}, x_n)}|f(x_n) - f(x_{n-1})|$ . We now sum over all  $x_i$ , use the definition of  $m$  and the triangle inequality to obtain that

$$\begin{aligned} \sum_{x \in \gamma} \rho(x) &\geq \sum_{i=0}^{n-1} \sqrt{c(x_i, x_{i+1})}|f(x_{i+1}) - f(x_i)| \\ (9) \qquad \qquad &\geq m|f(x_1) - f(x_0) + f(x_2) - f(x_1) + \dots + f(x_n) - f(x_{n-1})| \\ &\geq mg. \end{aligned}$$

Assertion (2) of the theorem now follows.

**3.5**

**Remark.** It is easy to check that Assertion (2) of the theorem will hold for a larger class of horizontal paths.

We now provide an upper bound for the product of the lengths of any shortest paths in the network.

**Lemma 3.6** *Let  $\mathcal{T}$  be a triangulation of a topological quadrilateral. Let  $\Gamma(\bar{F})$  be the associated network. Let  $f$  be the Dirichlet-Neumann boundary value function with  $g \neq 0$ . Let  $\rho$  be the gradient metric induced by  $f$ . Then for any  $\rho$  shortest curves  $\alpha, \beta$  in  $\Gamma(\bar{F})$  we have that*

$$\text{length}_\rho(\alpha)\text{length}_\rho(\beta) \leq l(|V|)I(f),$$

where  $l(|V|)$  is some constant which depends on  $|V|$ .

**Proof.** Let  $\alpha = (x_0, x_1, \dots, x_n)$  be a  $\rho$  shortest curve in  $\Gamma(\bar{F})$  connecting the vertex  $x_0$  to the vertex  $x_n$ . Then  $\text{length}_\rho(\alpha) = \sum_{x \in \alpha} \rho(x)$ . By the definition of  $\rho$  (Definition 3.3) we have for all  $x \in V$  that

$$\rho(x) = \left( \sum_{y \in \bar{F}} c(x, y)(f(x) - f(y))^2 \right)^{1/2} \leq \left( \sum_{(x, y) \in \bar{E}} c(x, y)(f(x) - f(y))^2 \right)^{1/2} = \sqrt{I(f)}.$$

It follows from Chapter 31 in [17] (with only minor changes needed in our setting) that  $n = O((|E| + |V|) \log |V|)$ . Since  $\Gamma(\bar{F})$  is planar we also have that  $|E| = O(|V|)$ . Hence we have that  $n = O(|V| \log |V|)$ . Therefore it follows that

$$\text{length}_\rho(\alpha) = O(|V| \log |V|) \sqrt{I(f)}.$$

The assertion of the lemma follows easily.

**3.6**

**Corollary 3.7** *Under the assumptions of the Theorem 3.5 and Lemma 3.6 we have that*

$$l(|V|)I(f) \geq \text{length}_\rho(\gamma^*)\text{length}_\rho(\gamma) \geq \frac{m}{M}I(f).$$

**Remark.** In the case that  $c(x, y) \equiv 1$ , it is easy to see that  $\frac{m}{M} = \frac{1}{\sqrt{k}}$ .

## 4 An example




Figure 3

With the triangulation above, let us solve the Dirichlet-Neumann boundary value (Definition 3.1) with  $c(x, y) \equiv 1$  and  $g = 1$ . By abuse of notation let us use the same letter to indicate both the vertex name and the value of the solution at this vertex. Simple calculations show the following.

1.  $(X, V, S, T, Y, L, U) = (1/2, 1/2, 3/4, 1/4, 1/4, 3/4, 1/2)$  is the solution.
2.  $[0, X, 1]$  is the shortest (proper) horizontal geodesic and its length, denoted by  $l_h$ , with respect to the gradient metric is equal to  $\sqrt{3/2} + \frac{\sqrt{3}}{2}$  ( $\sim 2.09077$ ),
3.  $[V, X, U]$  is the shortest (proper) vertical geodesic and its length, denoted by  $l_v$ , with respect to the gradient metric is equal to  $\frac{\sqrt{3}}{2}$  ( $\sim 0.866025$ ),
4. the energy,  $I$  of the solution equals  $3/2$ ,
5.  $\frac{l_c l_v}{I} - \frac{1}{\sqrt{8}} \sim 0.853553 > 0$ , and
6.  $\frac{8I}{l_c l_v} \sim 6.62742 > 1$ .

We conclude that the bounds provided by the remark following Corollary 3.7 are not sharp.

## References

- [1] L.V. Ahlfors and L. Sario, *Riemann Surfaces*, Princeton N.J., Princeton University Press, 1960.
- [2] E. Bendito, A. Carmona, A.M. Encinas, *Solving boundary value problems on networks using equilibrium measures*, J. of Func. Analysis, **171** (2000), 155-176.

- [3] E. Bendito, A. Carmona, A.M. Encinas, *Shortest Paths in Distance-regular Graphs*, Europ. J. Combinatorics, **21** (2000), 153-166.
- [4] E. Bendito, A. Carmona, A.M. Encinas, *Equilibrium measure, Poisson Kernel and Effective Resistance on Networks*, De Gruyter. Proceeding in Mathematics, (V. Kaimanovich, K. Schmidt, W. Woess ed.), **174** (2003), 363-376.
- [5] E. Bendito, A. Carmona, A.M. Encinas, *Difference schemes on uniform grids performed by general discrete operators*, Applied Numerical Mathematics, **50** (2004), 343-370.
- [6] I. Benjamini, O. Schramm, *Harmonic functions on planar and almost planar graphs and manifolds, via circle packings*, Invent. Math. **126** (1996), 565-587.
- [7] A.S. Besicovitch, *On two problems of Loewner*, J. London Math. Soc. **27** (1952), 141-144.
- [8] J. W. Cannon, *The combinatorial Riemann mapping theorem*, Acta Math. **173** (1994), 155 - 234.
- [9] J.W. Cannon, W.J. Floyd and W.R. Parry, *Squaring rectangles: the finite Riemann mapping theorem*, Contemporary Mathematics, vol. 169, Amer. Math. Soc., Providence, 1994, 133 - 212.
- [10] F.R. Chung, A. Grigořyan, S.T. Yau, *Upper bounds for eigenvalues of the discrete and continuous Laplace operators*, Adv. Math. **117** (1996), 165-178.
- [11] R. Duffin, *The extremal length of a network*, J. Math. Anal. Appl. **5**, 1962, 200-215.
- [12] B. Fuglede, *On the theory of potentials in locally compact spaces*, Acta. Math. **103** (1960), 139-215.
- [13] S. Hersensky, *Making a quadrilateral (or a ring) straight*, in preparation.
- [14] A. Marden and B. Rodin, *Extremal and conjugate extremal distance on open Riemann surfaces with applications to circular-radial slit mappings*, Acta. Math. **115** (1966), 237-269.
- [15] M. Ohtsuka, *Dirichlet problem, extremal length, and prime ends*, Van Nostrand Reinhold mathematical studies **22** (1970).
- [16] O. Schramm, *Square tilings with prescribed combinatorics*, Israel Jour. of Math. **84** (1993), 97-118.
- [17] R. Sedgewick, *Algorithms*, 2nd edition, Addison-Wesley, 1988.
- [18] P.M. Sordani, *Potential Theory on Infinite Networks*, Lecture Notes in Math. **1590**, Springer, Berlin, 1994.
- [19] W. Woess, *Random walks on infinite graphs and groups*, Cambridge tracts in mathematics, **138**, Cambridge ; New York : Cambridge University Press, 2000.

University of Georgia  
 Dept. of Mathematics  
 Athens, GA 30602  
 saarh@math.uga.edu